# REMARKS ON PROJECTIVE NORMALITY FOR CERTAIN CALABI-YAU AND HYPERKÄHLER VARIETIES

JAYAN MUKHERJEE AND DEBADITYA RAYCHAUDHURY

ABSTRACT. We prove some results on effective very ampleness and projective normality for some varieties with trivial canonical bundle. In the first part we prove an effective projective normality result for an ample line bundle on regular smooth four-folds with trivial canonical bundle. More precisely we show that for a regular smooth fourfold with trivial canonical bundle,  $A^{\otimes 15}$  is projectively normal for *A* ample. In the second part we emphasize on the projective normality of multiples of ample and globally generated line bundles on certain classes of known examples (upto deformation) of projective hyperkähler varieties. As a corollary we show that excepting two extremal cases in dimensions 4 and 6, a general curve section of any ample and globally generated linear system on the above mentioned examples is non-hyperelliptic.

### INTRODUCTION

In this article, we prove some effective projective normality results for multiples of line bundles with some positivity conditions on Calabi-Yau and hyperkähler varieties. These are varieties with trivial canonical bundles that are higher dimensional analogues of *K*3 surfaces. We recall some results on syzygies for varieties with trivial canonical bundle below and state our main result for regular four-folds with trivial canonical bundles. Next we will concentrate on a special class of even dimensional such varieties (regular with trivial canonical bundle), namely hyperkähler varieties.

0.1. **Regular varieties with trivial canonical bundle.** Geometry of linear series on varieties X with trivial canonical bundle  $K_X$  (i.e.  $K_X = 0$ ) is a topic that has motivated a lot of research. The question of what multiple of an ample line bundle on a K-trivial variety is very ample and projectively normal was extensively studied by many mathematicians including Gallego, Oguiso, Peternell, Purnaprajna, Saint-Donat (see [11], [29], [30]). Recall that a K3 surface is defined as a smooth projective surface S with  $K_S = 0$  and  $H^1(\mathcal{O}_S) = 0$ . The following theorem is due to Saint-Donat (see [30]).

**Theorem 0.1.** Let *S* be a smooth projective K3 surface and let *B* be an ample line bundle on *S*. Then  $B^{\otimes n}$  is very ample (in fact projectively normal) for  $n \ge 3$ .

Gallego and Purnaprajna generalized Saint-Donat's result on projective normality for smooth, projective, regular ( $H^1(\mathcal{O}_X) = 0$ ) three-fold X with trivial canonical bundle (see [11], Corollary 1.10). They showed that for an ample line bundle B on X,  $B^{\otimes n}$  is projectively normal for n = 8 and  $n \ge 10$ . Moreover, if  $B^3 > 1$  then  $B^{\otimes n}$  is projectively normal if n = 6 and  $n \ge 8$ . In order to prove this theorem, Gallego and Purnaprajna studies the case when a regular, three-fold with trivial canonical bundle maps onto a variety of minimal degree by a complete linear series of an ample and globally generated line bundle. They gave a classification theorem for such situations and proved

the projective normality result as a consequence by applying a theorem of Green (see Theorem 1.7). Varieties that appear as covers of varieties of minimal degree play an important role in the geometry of algebraic varieties. They are extremal cases in a variety of situations from algebraic curves to higher dimensional varieties (see [11], [12], [13], [16]).

In this article, we prove an analogue of the classification theorem of Gallego and Purnaprajna where we study the situation when a smooth regular four-fold X with trivial canonical bundle maps to a variety of minimal degree by the complete linear system of an ample and globally generated line bundle B (see Theorem 2.3) and provide upper bounds to the degree of such morphisms. As a consequence of the classification theorem, Fujita's base point freeness conjecture that has been proved in dimension four by Kawamata (see [18]) and Green's theorem (c.f. Theorem 1.7), we prove the following effective projective normality result (see Theorem 2.4) on regular fourfolds with trivial canonical bundle.

**Theorem A.** *Let X be a four-fold with trivial canonical bundle and let A be an ample line bundle on X. then,* 

- (1) *nA* is very ample and embeds X as a projectively normal variety for  $n \ge 16$ .
- (2) If  $H^1(\mathcal{O}_X) = 0$  then nA is very ample and embeds X as a projectively normal variety for  $n \ge 15$ .

This theorem can be thought of as a higher dimensional analogue of Theorem 0.1 and Corollary 1.10, [11]. Note that standard methods of Castelnuovo-Mumford regularity (see Lemma 1.6) and Theorem 1.3, [11] yields in the situation above that  $A^{\otimes n}$  satisfies projective normality for  $n \ge 21$ .

0.2. Calabi-Yau and Hyperkähler varieties. The definition of K3 surface, i.e. a smooth regular surface S with trivial canonical bundle is equivalent to having a holomorphic symplectic form on S. However in higher dimensions these two notions do not coincide which is clear from the fact that existence of a holomorphic symplectic form on a Kähler manifold demands that its dimension is even whereas there are examples of smooth projective algebraic varieties in odd dimensions with trivial canonical bundle and  $H^1(\mathcal{O}_X) = 0$ , for example smooth hypersurfaces of degree n + 1 in  $\mathbb{P}^n$ . So essentially we can have two different kinds of generalizations of a K3 surface. We give the precise definitions of Calabi-Yau and hyperkähler varieties below.

**Definition 0.2.** A compact Kähler manifold M of dimension  $n \ge 3$  is called Calabi-Yau if it has trivial canonical bundle and the hodge numbers  $h^{p,0}(M)$  vanish for all 0 .

With this definition, Calabi-Yau manifolds are neccessarily projective. However the following definition of hyperkähler manifolds does not imply projectivity in general.

**Definition 0.3.** A compact Kähler manifold M is called hyperkähler if it is simply connected and its space of global holomorphic two forms is spanned by a symplectic form.

The decomposition theorem of Bogomolov (see [2]) says, any complex manifold with trivial first Chern class admits a finite étale cover isomorphic to a product of complex tori, Calabi-Yau manifolds and hyperkähler manifolds. Thus, these spaces can be thought as the "building blocks" for manifolds with trivial first Chern class.

The theorem of Saint-Donat that we stated (see Theorem 0.1) deals with ample line bundles.

In the same paper (see [30]), Saint-Donat proves the following theorem for ample and globally generated line bundles on K3 surfaces.

**Theorem 0.4.** Let *S* be a smooth projective K3 surface and let *B* be an ample and base point free line bundle on *S*. Then,

- (1)  $B^{\otimes 2}$  is very ample and  $|B^{\otimes 2}|$  embeds *S* as a projectively normal variety unless the morphism given by the complete linear system |B| maps *S*, 2 : 1 onto  $\mathbb{P}^2$ .
- (2) *B* is very ample and |B| embeds *S* as a projectively normal variety unless the morphism given by the complete linear system |B| maps *S*, 2 : 1 onto  $\mathbb{P}^2$  or to a variety of minimal degree.

Gallego and Purnaprajna provides a generalization of this theorem for regular three-folds X with trivial canonical bundle in [11] in which they proved for an ample and globally generated line bundle B,  $B^{\otimes 3}$  is projectively normal unless the morphism  $\varphi_B$  induced by the complete linear series of B maps X, 2-1 onto  $\mathbb{P}^3$ . Moreover, they showed that  $B^{\otimes 2}$  is projectively normal unless  $\varphi_B$  maps X, 2-1 onto  $\mathbb{P}^3$  or to a variety of minimal degree  $\geq 2$ . They also proved that  $B^{\otimes 4}$  is projectively normal on smooth, projective, regular, four-folds with trivial canonical bundle when the morphism  $\varphi_B$  associated to the complete linear series of an ample and globally generated line bundle B is birational onto its image and  $h^0(B) \geq 7$  (see Theorem 1.11, [11]). Recently Niu proved an analogue of Theorem 0.4 in dimension four. In fact he proved a general result for smooth, projective, regular,  $K_X$ -trivial varieties in all dimensions (see [26]) with an additional assumption of  $H^2(\mathcal{O}_X) = 0$  which in dimensions 2 and 3 recovers, and in dimension 4 generalizes the results of Saint-Donat and Gallego-Purnaprajna.

We see that it is a natural question to ask whether and to what extent these theorems generalize to the other class of higher dimensional analogues of K3 surfaces, namely hyperkähler varieties. Recall that there are many families of examples for Calabi-Yau varieties but only few classes of examples for hyperkähler varieties are known upto deformation. Hilbert scheme of *n* points on a K3 surface (we will denote them by  $K3^{[n]}$ ), generalized Kummer varieties (we will denote them by  $K^n(T)$ ) and two examples in dimension six and ten given by O'Grady (we will denote it by  $\mathcal{M}_6$  and  $\mathcal{M}_{10}$  respectively) are the only known classes of examples upto deformation. Our main theorem for hyperkähler varieties is the following.

**Theorem B.** Let X be a projective hyperkähler variety of dimension  $2n \ge 4$  that is deformation equivalent to  $K3^{[n]}$  or  $K^n(T)$  or  $\mathcal{M}_6$ . Let B be an ample and globally generated line bundle on X. Then the following happens.

(1)  $B^{\otimes l}$  is projectively normal for  $l \geq 2n$ .

(2)  $B^{\otimes 2n-1}$  is projectively normal unless:

- (a) n = 2,  $X = K3^{[2]}$  and  $\varphi_B$  maps X onto a quadric (possibly singular) inside  $\mathbb{P}^5$ . In this case  $q_X(B) = 2$ ,  $deg(\varphi_B) = 6$ , or
- (b) n = 3,  $X = K3^{[3]}$  and  $\varphi_B$  maps X onto a variety of minimal degree inside  $\mathbb{P}^9$  which is obtained by taking cones over the Veronese embedding of  $\mathbb{P}^2$  inside  $\mathbb{P}^5$ . In this case  $q_X(B) = 2$ ,  $deg(\varphi_B) = 30$ .

Hence if X is as above and B does not satisfy cases 2(a) or 2(b) then a general curve section of |B| is non-hyperelliptic.

### MUKHERJEE AND RAYCHAUDHURY

As before, the study of the situation when X maps onto a variety of minimal degree by the complete linear series |B| is the main ingredient of our proof. We also use two key characteristics of a hyperkähler variety which are the Riemann-Roch expression that comes from the existence of a primitive integral quadratic form on the second integral cohomology group of the variety and Matsushita's theorem on fibre space structure of a hyperkähler manifold (see Theorem 1.15).

Now we give the structure of our paper. In Section 2, we briefly recall the main theorems and observations that we will use to prove our main results. Section 3 and Section 4 deals with the proof of Theorem A and B respectively.

Acknowledgements. We want to thank our advisor Prof. B.P. Purnaprajna for his encouragement, support and guidance without which this work would have never been possible.

### 1. PRELIMINARIES AND NOTATIONS

Throughout this article, X will always denote a smooth, projective variety over  $\mathbb{C}$ . K or  $K_X$  will denote its canonical bundle. We will use the multiplicative and the additive notation of line bundles interchangeably. Thus, for a line bundle L,  $L^{\otimes r}$  and rL are the same. We have used the notation  $L^{-r}$  for  $(L^*)^{\otimes r}$ . We will use  $L^r$  to denote the intersection product.

1.1. **Projectve normality.** For a globally generated line bundle L on a smooth projective variety X, we have the following short exact sequence.

(1.1) 
$$0 \to M_L \to H^0(L) \otimes \mathscr{O}_X \to L \to 0.$$

First we recall the definition of projective normality and the  $N_p$  property.

**Definition 1.1.** *Let L be a very ample line bundle on a variety X. Let the following be the minimal graded free resolution of the coordinate ring R of the embedding of X induced by L* 

 $0 \to F_n \to F_{n-1} \to \cdots \to F_0 \to R \to 0.$ 

Let  $\mathscr{I}_X$  be the ideal sheaf of the embedding.

- (1) L satisfies the property  $N_0$  (or embeds X as a projectively normal variety) if R is normal.
- (2) L satisfies the property  $N_1$  (or is normally presented) if in addition  $\mathscr{I}_X$  is generated by quadrics.
- (3) L satisfies the property  $N_p$  if in addition to satisfying the property  $N_1$ , the resolution is linear from the second step until the p-th step.

We have the following necessary and sufficient condition for the  $N_p$  property of an ample and base point free line bundle on X. In this article, we will only deal with the case when p = 0.

**Theorem 1.2.** Let *L* be an ample, globally generated line bundle on *X*. Suppose the cohomology group  $H^1(\bigwedge^{p'+1} M_L \otimes L^{\otimes k})$  vanishes for all  $0 \le p' \le p$  and for all  $k \ge 1$ , then *L* satisfies the property  $N_p$ . If in addition  $H^1(L^{\otimes r}) = 0$  for all  $r \ge 1$ , then the above is a necessary and sufficient condition for *L* to satisfy  $N_p$ .

We have made use of the following observation of Gallego and Purnaprajna (see for instance [11]) to show the surjections of certain multiplication maps.

**Observation 1.3.** Let E and  $L_1, L_2, \ldots, L_r$  be coherent sheaves on a variety X. Consider the map  $H^0(E) \otimes H^0(L_1 \otimes L_2 \otimes \cdots \otimes L_r) \xrightarrow{\psi} H^0(E \otimes L_1 \otimes \cdots \otimes L_r)$  and the following maps

$$H^{0}(E) \otimes H^{0}(L_{1}) \xrightarrow{\alpha_{1}} H^{0}(E \otimes L_{1}),$$
$$H^{0}(E \otimes L_{1}) \otimes H^{0}(L_{2}) \xrightarrow{\alpha_{2}} H^{0}(E \otimes L_{1} \otimes L_{2}),$$

$$H^{0}(E \otimes L_{1} \otimes \cdots \otimes L_{r-1}) \otimes H^{0}(L_{r}) \xrightarrow{\alpha_{r}} H^{0}(E \otimes L_{1} \otimes \ldots \otimes L_{r}).$$

If  $\alpha_1, \alpha_2, \ldots, \alpha_r$  are surjective then  $\psi$  is also surjective.

The technique we use to show projective normalty of an ample and globally generated line bundle on a variety is to use Koszul resolution to restrict the bundle on a smooth curve section and then showing the surjectivity of an appropriate multiplication map. It is worth mentioning that Koszul resolution is the special case of a particular complex, known as Skoda complex (see [20]).

**Definition 1.4.** *Let X be a smooth projective variety of dimension*  $n \ge 2$ *. Let L be an ample and globally generated line bundle on X.* 

- (1) Take n 1 general sections  $s_1, \ldots, s_{n-1}$  of  $H^0(L)$  so the intersection of the divisor of zeroes  $L_i = (s_i)_0$  is a nonsingular projective curve C, that is  $C = L_1 \cap \cdots \cap L_{n-1}$ .
- (2) Let  $\mathscr{I}$  be the ideal sheaf of C and let  $W = span\{s_1, \ldots, s_{n-1}\} \subseteq H^0(L)$  be the subspace spanned by  $s_i$ . Note that  $W \subseteq H^0(L \otimes \mathscr{I})$ . The Koszul resolution  $\mathbf{I}_1$  is the following exact sequence,

$$0 \to \bigwedge^{n-1} W \otimes L^{-(n-1)} \to \cdots \to \bigwedge^2 W \otimes L^{-2} \to W \otimes L^{-1} \to \mathscr{I} \to 0.$$

Once we boil down our problem to a problem on curve, we use the following two results. The first one is a result of Green (see [13]).

**Lemma 1.5.** Let C be a smooth, irreducible curve. Let L and M be line bundles on C. Let W be a base point free linear subsystem of  $H^0(C, L)$ . Then the multiplication map  $W \otimes H^0(M) \to H^0(L \otimes M)$  is surjective if  $h^1(M \otimes L^{-1}) \leq \dim(W) - 2$ .

The second one is known as Castelnuovo-Mumford lemma (see [24]). It is one of the main tools that has been used extensively in studying the syzygies of a variety embedded in a projective space.

**Lemma 1.6.** Let *L* be a base point free line bundle on a variety *X* and let  $\mathscr{F}$  be a coherent sheaf on *X*. If  $H^i(\mathscr{F} \otimes L^{-i}) = 0$  for all  $i \ge 1$  then the multiplication map  $H^0(\mathscr{F} \otimes L^{\otimes i}) \otimes H^0(L) \to H^0(\mathscr{F} \otimes L^{\otimes i+1})$  surjects for all  $i \ge 0$ .

One can obtain effective projective normality results using the above theorems and lemmas. In order to improve the bounds one can use the following version of a result of Green (see [12], p.1089, (3)).

**Theorem 1.7.** Let X be a regular projective variety of dimension n for which the canonical bundle K is ample and base point free. Moreover, assume  $h^0(K) \ge n + 2$ . Let  $\varphi_K$  be the morphism induced by the complete linear series |K|. If  $\varphi_K(X)$  is not a variety of minimal degree then the multiplication map  $H^0(K) \otimes H^0(K^{\otimes l}) \to H^0(K^{\otimes l+1})$  surjects for  $l \ge n$ .

The following corollary is the precise version of what we will use in the subsequent sections.

**Corollary 1.8.** Let X be a regular variety of dimension  $n \ge 3$  with trivial canonical bundle. Let L be an ample and globally generated line bundle on X and  $\varphi_L$  be the morphism induced by the complete linear series |L|. Assume  $\varphi_L$  does not map X onto a variety of minimal degree. Then  $L^{\otimes n-1}$  is projectively normal.

*Proof.* We use Theorem 1.2 and Observation 1.3 to notice that to prove the projective normality of  $L^{\otimes n-1}$ , it is enough to show the surjectivity of the map  $H^0(lL) \otimes H^0(L) \longrightarrow H^0((l+1)L)$  for  $l \ge n-1$ . We just prove the case for l = n - 1 here, the rest follow similarly.

Choose a smooth threefold section T of the ample and base point free line bundle L. We have the following commutative diagram,

where  $K_T = B|_T$  (by adjunction) denotes the canonical bundle of *T*. Since the leftmost map surjects, the middle vertical map surjects if and only if rightmost vertical map surjects. By Kodaira vanishing we have that  $H^1((n-2)L) = 0$  and hence we have a surjection,

$$H^0((n-1)L) \to H^0((n-1)L|_T) = H^0((n-1)K_T).$$

Hence the rightmost vertical map surjects if and only if we have the surjection of

$$H^0((n-1)K_T) \otimes H^0(K_T) \longrightarrow H^0(nK_T).$$

We notice that *T* is a smooth irreducble regular variety of general type and hence by Theorem 1.7, the above map surjects unless *T* is mapped to a variety of minimal degree by the complete linear series of  $L|_T$ . But the latter is equivalent to saying that *X* is mapped by the complete linear series of |L| to a variety of minimal degree.

We briefly recall some basic definitions and results concerning varieties of minimal degree.

## 1.2. Varieties of minimal degree. For any non-degenerate variety $X \subset \mathbb{P}^r$ , we have the inequality

$$deg(X) \ge 1 + codim(X).$$

A variety  $X \subset \mathbb{P}^r$  is said to be a *variety of minimal degree* if it satisfies the equality deg(X) = 1 + codim(X). If codim(X) = 1 then of course the variety X is a quadric hypersurface. The following theorem of Eisenbud and Harris (see [6]) provides a complete classification of the varieties of minimal degree.

**Theorem 1.9.** Let  $X \subset \mathbb{P}^r$  is a variety of minimal degree. Then X is a cone over a smooth such variety. Moreover, if X is smooth and codim(X) > 1 then  $X \subset \mathbb{P}^r$  is either a rational normal scroll or a Veronese surface  $\mathbb{P}^2 \subset \mathbb{P}^5$ .

Recall that a rational normal scroll  $X \subset \mathbb{P}^r$  of dimension *n* is the image of a projective bundle  $\pi : \mathbb{P}(\mathscr{E}) \to \mathbb{P}^1$  through the morphism goven by the tautological line bundle  $\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)$  where the vector bundle  $\mathscr{E} = \mathscr{O}(a_1) \oplus \cdots \oplus \mathscr{O}(a_n)$  satisfies  $0 \le a_1 \le a_2 \le \cdots \le a_n$  and  $deg(X) = \sum a_i$ .

If  $a_1 = a_2 = \cdots = a_l = 0$  for some 0 < l < n then X singular and it is a cone over a smooth rational normal scroll. The vertex or singular locus V of this cone has dimension l - 1 and let  $X_S = X \setminus V$  be the smooth part of X. Moreover, X is normal and  $\tilde{X} = \mathbb{P}(\mathscr{E}) \to X$  is a rational resolution of singularity which is called the *canonical resolution* of the rational normal scroll X.

Let X be a rational normal scroll. Let H be the class of a hyperplane section and R be the class of a general linear subspace of codimension one. We note the following fact:

**Lemma 1.10.** If codim(V, X) = 2 then  $H \sim deg(X)R$ .

*Proof.* See [8], Corollary 2.2 (2).

We prove a lemma here that we will use in the subsequent sections to prove our main theorems.

**Lemma 1.11.** Let X be a smooth projective variety and let L be an ample and globally generated line bundle on X. Let  $\varphi_L$  be the morphism induced by the complete linear series L. Assume  $\varphi_L$  maps X onto a singular rational normal scroll Y with vertex V. Then codim(V, Y) = 2 that is Y is obtained by taking cones over a rational normal curve.

*Proof.* Suppose  $\tilde{Y} = \mathbb{P}(\mathscr{E}) \to Y$  is the canonical resolution. If the codimension of the singular locus is > 2, then the canonical resolution is a small resolution (see [8], Proposition 2.1) and hence *X* could be obtained by contracting subschemes of  $(X \times_Y \tilde{Y})_{red}$  of codimension greater than one, which contradicts the factoriality of *X*. The assertion follows since  $codim(V, Y) \neq 1$  as *Y* is normal.  $\Box$ 

1.3. Hyperkähler varieties. Recall that a hyperkähler variety X is a compact Kähler manifold that is simply connected, projective and its space of global holomorphic two forms is spanned by a symplectic form. The symplectic form ensures that  $K_X$  is trivial and dim(X) is even. It is also known that  $\chi(\mathscr{O}_X) = n + 1$  and the following are the values of  $h^p(X, \mathscr{O}_X)$ ,

$$h^{p}(X, \mathscr{O}_{X}) = \begin{cases} 1 & p \text{ is even,} \\ 0 & p \text{ is odd.} \end{cases}$$

Only a few classes of examples of hyperkähler varieties are known. Beauville first gave examples of two distinct deformation classes of compact hyperkähler manifolds in all even dimensions greater than or equal to 2 (see [1]). The first example is the Hilbert scheme  $S^{[n]}$  of length *n* subschemes on a *K*3 surface *S*. The second one is the generalized Kummer variety  $K^n(T)$  which is the fibre over the 0 of an Abelian variety *T* under the morphism  $\phi \circ \psi$  (see the diagram below)

$$T^{[n+1]} \xrightarrow{\psi} T^{(n+1)} \xrightarrow{\phi} T$$

where  $T^{[n+1]}$  Hilbert scheme of length n + 1 subschemes on the Abelian variety T,  $T^{(n+1)}$  is the symmetric product,  $\psi$  is the Hilbert chow morphism and  $\phi$  is the addition on T. Two other distinct deformation classes of hyperkahler manifolds  $\mathcal{M}_6$  and  $\mathcal{M}_{10}$  are given by O'Grady in dimensions 6 and 10 respectively which appear as desingularizations of certain modulii spaces of sheaves over symplectic surfaces (see [27], [28]). All other known examples are deformation equivalent to one of these.

We start by the following theorem of Beauville and Fujiki (see [1] and [9]).

**Theorem 1.12.** Let X be a hyperkähler variety of dimension 2n. There exists a quadratic form  $q_X$ :  $H^2(X, \mathbb{C}) \to \mathbb{C}$  and a positive constant  $c_X \in \mathbb{Q}_+$  such that for all  $\alpha$  in  $H^2(X, \mathbb{C})$ ,  $\int_X \alpha^{2n} = c_X \cdot q_X(\alpha)^n$ . The above equation determines  $c_X$  and  $q_X$  uniquely if one assumes the following two conditions.

- (I)  $q_X$  is a primitive integral quadratic form on  $H^2(X, \mathbb{Z})$ ;
- (II)  $q_X(\sigma, \bar{\sigma}) > 0$  for all  $0 \neq \sigma \in H^{2,0}(X)$ .

Here  $q_X$  and  $c_X$  are called the Beauville form and Fujiki constant respectively.

The Beauville form and Fujiki constants are fundamental invariants of a hyperkähler variety. They play an important role in determining the intersections on X as the following theorem shows (See [9], [14]).

**Theorem 1.13.** Let X be a hyperkähler variety of dimension 2n. Assume that  $\alpha \in H^{4j}(X, \mathbb{C})$  of type (2j, 2j) on all small deformations of X. Then there exists a constant  $C(\alpha) \in \mathbb{C}$  depending on  $\alpha$  such that  $\int_{X} \alpha \cdot \beta^{2n-2j} = C(\alpha) \cdot q_X(\beta)^{n-j}$  for all  $\beta \in H^2(X, \mathbb{C})$ .

As a consequence of the theorem above, we get the following form of the Riemann-Roch formula for an line bundle L on a hyperkähler variety of dimension 2n (see [17]),

(1.2) 
$$\chi(X,L) = \sum_{i=0}^{n} \frac{a_i}{(2i)!} q_X(c_1(L))^i$$

where  $a_i = C(td_{2n-2i}(X))$ . Here  $a_i$ 's are constants depending only on the topology of X.

Elingsrad-Gottsche-Lehn computes the rational constants of the Riemann roch expression for hyperkähler manifolds of deformation type  $K3^{[n]}$  (See [7]) and Britze-Nieper computes the same for generalized Kummer varieties  $K^n(T)$  of dimension 2n (see [3]). If X is of  $K3^{[n]}$  type we have,

(1.3) 
$$\chi(L) = \begin{pmatrix} \frac{1}{2}q(L) + n + 1 \\ n \end{pmatrix} \quad and \quad c_X = \frac{(2n)!}{n!2^n}$$

For a generalized Kummer variety of dimension 2n we have the following Riemann-Roch formula,

(1.4) 
$$\chi(L) = (n+1) \binom{\frac{1}{2}q(L)+n}{n} \quad and \quad c_X = (n+1) \frac{(2n)!}{n!2^n}$$

The Riemann-Roch formula and Fujiki constant for  $\mathcal{M}_6$  are the same as that of  $K^3(T)$ .

We will use Matsushita's theorem on fibre space structure of hyperkähler varieties. We recall the the definition and the main theorem.

**Definition 1.14.** Let X be an algebraic variety. A fibre space structure of X is a proper surjective morphism  $f : X \to S$  that satisfies the following two conditions:

- (1) *X* and *S* are normal varieties with  $0 < \dim(S) < \dim(X)$ .
- (2) A general fibre of f is connected.

The result that we will use in Section 4 is the following (see [21], Theorem 2, (3)).

**Theorem 1.15.** Let  $f : X \to B$  is a fibre space structure on a projective hyperkähler variety X of dimension 2n with projective base B. Then dim(B) = n.

### 2. Proof of Theorem A

The main aim of this section is to prove results on effective very ampleness and projective normality on a four dimensional variety with trivial canonical bundle. We start with a general statement on projective normality and normal presentation. **Lemma 2.1.** Let X be a smooth, projective n-fold with trivial canonical bundle. Let B be an ample and base point free line bundle on X. Let  $h^0(B) \ge n + 2$ . Then lB satisfies the property  $N_0$  for all  $l \ge n$ . Moreover, if X is Calabi-Yau, then lB satisfies the property  $N_1$  for all  $l \ge n$ .

*Proof.* Follows immediately from Theorem 2.3 and Theorem 3.4 of [23].

Now we want to find out what multiple of an ample line bundle is very ample on a four dimensional variety with trivial canonical bundle. We will use the Fujita freeness on four folds that has been proved by Kawamata in [18]. We begin with a lemma.

**Lemma 2.2.** Let X be a fourfold with trivial canonical bundle. Let A be an ample line bundle and let B = nA for  $n \ge 5$ . Then the multiplication map  $H^0(3B+kA) \otimes H^0(B) \rightarrow H^0(4B+kA)$  is surjective for  $k \ge 1$ .

*Proof.* Note that *B* is base point free by Kawamata's proof of Fujita's base point freeness theorem on fourfolds (see [18]). We prove the statement for k = 1. For k > 1 the proof is exactly the same.

Let *C* be a smooth and irreducible curve section of the linear system |B| and let  $\mathscr{I}$  be the ideal sheaf of *C* in *X*. We have the following commutative diagram with the two horizontal rows exact. Here  $\mathscr{I}$  is the ideal sheaf of *C* in *X*, *V* is the cokernel of the map  $H^0(B \otimes \mathscr{I}) \to H^0(B)$ .

$$\begin{array}{cccc} 0 \rightarrow H^{0}(B \otimes \mathscr{I}) \otimes H^{0}(3B + A) \rightarrow H^{0}(B) \otimes H^{0}(3B + A) \rightarrow V \otimes H^{0}(3B + A) \rightarrow 0 \\ & & \downarrow & & \downarrow \\ 0 \longrightarrow H^{0}((4B + A) \otimes \mathscr{I}) \longrightarrow H^{0}((4B + A)) \longrightarrow H^{0}(4B + A|_{C}) \longrightarrow 0 \end{array}$$

Now we claim that the leftmost vertical map is surjective. Consider the Koszul resolution,

(2.1) 
$$0 \to \bigwedge^{3} W \otimes B^{-3} \to \bigwedge^{2} W \otimes B^{-2} \to W \otimes B^{*} \to \mathscr{I} \to 0.$$

Tensor it with 4B + A to get the following,

$$(2.2) \qquad 0 \to \bigwedge^{3} W \otimes (B+A) \xrightarrow{f_{3}} \bigwedge^{2} W \otimes (2B+A) \xrightarrow{f_{2}} W \otimes (3B+A) \xrightarrow{f_{1}} (4B+A) \otimes \mathscr{I} \to 0$$

That gives us two short exact sequences.

(2.3) 
$$0 \to Ker(f_1) \to W \otimes (3B+A) \xrightarrow{f_1} (4B+A) \otimes \mathscr{I} \to 0,$$

(2.4) 
$$0 \to \bigwedge^{3} W \otimes (B+A) \xrightarrow{f_{3}} \bigwedge^{2} W \otimes (2B+A) \xrightarrow{f_{2}} Ker(f_{1}) \to 0.$$

Taking long exact sequence of cohomology in the second sequence we get the following,

(2.5) 
$$\bigwedge^{2} W \otimes H^{1}(2B+A) \to H^{1}(Ker(f_{1})) \to \bigwedge^{3} W \otimes H^{2}(B+A).$$

Hence  $H^1(Ker(f_1)) = 0$  since the other terms of the exact sequence vanish by Kodaira Vanishing. The long exact sequence of cohomology associated to the first sequence is the following,

(2.6) 
$$W \otimes H^0(3B + A) \to H^0((4B + A) \otimes \mathscr{I})) \to H^1(Ker(f_1)).$$

We showed that the last term is zero, thus  $W \otimes H^0(3B + A) \to H^0((4B + A) \otimes \mathscr{I}))$  surjects. Consequently,  $H^0(B \otimes \mathscr{I}) \otimes H^0(3B + A) \to H^0((4B + A) \otimes \mathscr{I})$  surjects since  $W \subseteq H^0(B \otimes \mathscr{I})$ .

In order to prove the lemma we are left to show that  $V \otimes H^0(3B + A) \to H^0((4B + A)|_C)$  surjects. Since we have the surjection of  $H^0(3B + A) \to H^0((3B + A)|_C)$ , it is enough to show the surjection of  $V \otimes H^0(3B + A|_C) \to H^0((4B + A)|_C)$ . Thus, using Lemma 1.5 we need to prove that

$$h^1(2B+A|_C) \leq dimV-2$$

To prove this inequality, first we tensor the Koszul resolution by B,

(2.7) 
$$0 \to \bigwedge^{3} W \otimes B^{-2} \xrightarrow{f_{3}} \bigwedge^{2} W \otimes B^{*} \xrightarrow{f_{2}} W \otimes \mathscr{O}_{X} \xrightarrow{f_{1}} B \otimes \mathscr{I} \to 0.$$

As before, we end up getting two short exact sequences,

(2.8) 
$$0 \to Ker(f_1) \to W \otimes \mathscr{O}_X \xrightarrow{f_1} B \otimes \mathscr{I} \to 0$$

(2.9) 
$$0 \to \bigwedge^{3} W \otimes B^{-2} \xrightarrow{f_{3}} \bigwedge^{2} W \otimes B^{*} \xrightarrow{f_{2}} Ker(f_{1}) \to 0$$

The long exact sequence of cohomology associated to the second sequence gives,

(2.10) 
$$\bigwedge^{2} W \otimes H^{0}(B^{*}) \to H^{0}(Ker(f_{1})) \to \bigwedge^{3} W \otimes H^{1}(B^{-2})$$

Consequently,  $H^0(Ker(f_1)) = 0$  since  $H^1(B^{-2}) = 0$  by Kodaira vanishing and  $H^0(B^*) = 0$ . Taking cohomology once more we have the following exact sequence,

(2.11) 
$$\bigwedge^{2} W \otimes H^{1}(B^{*}) \to H^{1}(Ker(f_{1})) \to \bigwedge^{3} W \otimes H^{2}(B^{-2}).$$

Hence  $H^1(Ker(f_1)) = 0$  since the other terms of the exact sequence vanish by Kodaira Vanishing. The long exact sequence of cohomology associated to the first sequence is the following.

(2.12) 
$$H^{0}(Ker(f_{1})) \to W \otimes H^{0}(\mathscr{O}_{X}) \to H^{0}(B \otimes \mathscr{I})) \to H^{1}(Ker(f_{1}))$$

But the first and last terms are zero by Kodaira Vanishing and hence  $h^0(B \otimes \mathscr{I}) = dimW \le 3$ . Thus we obtain the inequality  $dimV - 2 \ge h^0(B) - 5$ .

On the other hand the canonical bundle of *C* is given by  $3B|_C$ . Applying Serre-Duality it is enough to prove that  $h^0(B - A) \le h^0(B) - 5$  i.e.  $h^0((n - 1)A) \le h^0(nA) - 5$ .

Applying Riemann Roch for nA and (n - 1)A and subtracting the equations we obtain,

(2.13) 
$$h^{0}(nA) - h^{0}((n-1)A) = \frac{n^{4} - (n-1)^{4}}{24}A^{4} + \frac{n^{2} - (n-1)^{2}}{24}A^{2} \cdot c_{2}.$$

The result of Miyaoka (see [22]) shows that  $A^2 \cdot c_2 \ge 0$  which gives,

(2.14) 
$$h^{0}(nA) - h^{0}((n-1)A) \ge \frac{n^{4} - (n-1)^{4}}{24}A^{4} \ge 5 \text{ if } n \ge 5$$

and that concludes the proof.

Now we give a classification theorem in which we classify the varieties which come as an image of a regular fourfold with trivial canonical bundle by an ample, globally generated line bundle with an additional property of being a variety of minimal degree.

**Theorem 2.3.** Let X be a regular four-fold with trivial canonical bundle. Let  $\varphi$  be the morphism induced by the complete linear series of an ample and base point free line bundle B on X with  $h^0(B) = r + 1$  and let d be the degree of  $\varphi$ . If  $\varphi$  maps X to a variety of minimal degree Y then,

$$d \le \frac{24(r-1)}{r-3}.$$

(a) Assume Y is smooth. Then one of the following happens:

- (1)  $Y = \mathbb{P}^4$ .
- (2) *Y* is a smooth quadric hypersurface in  $\mathbb{P}^5$ .
- (3) *Y* is a smooth rational normal scroll of dimension 4 in  $\mathbb{P}^6$  or  $\mathbb{P}^7$  and *X* is fibered over  $\mathbb{P}^1$ . Moreover, the general fibre is a smooth threefold *G* with  $K_G = 0$  and the degree *d* satisfies the following bounds;

$$2 \le d \le \min \Big\{ 6h^0(B|_G), \frac{24(r-1)}{r-3} \Big\}.$$

*If in addition G is regular we have the following;* 

$$2h^{0}(B|_{G}) - 6 \le d \le \min\left\{6(h^{0}(B|_{G}) - 1), \frac{24(r-1)}{r-3}\right\}, \text{ if } d \text{ is even and}$$
$$2h^{0}(B|_{G}) - 5 \le d \le \min\left\{6(h^{0}(B|_{G}) - 1), \frac{24(r-1)}{r-3}\right\}, \text{ if } d \text{ is odd.}$$

(4) *Y* is a smooth rational normal scroll in  $\mathbb{P}^r$  for  $r \ge 8$  and *X* is fibered over  $\mathbb{P}^1$  and the general fibre is a three-fold *G* with  $K_G = 0$  and the degree *d* of  $\varphi_B$  satisfies  $2 \le d \le 18$ .

(b) Assume Y is singular. Then one of the following happens:

- (1) *Y* is a singular quadric hypersurface.
- (2) *Y* is a singular four-fold which is either a triple cone over a rational normal curve in  $\mathbb{P}^r$  where  $6 \le r \le 8$  or a double cone over the Veronese surface in  $\mathbb{P}^5$ .

Proof. We first prove the inequality. Using Riemann-Roch we can see that,

(2.15) 
$$h^{0}(B) = \frac{1}{24}B^{4} + \frac{1}{24}B^{2} \cdot c_{2} + 2.$$

and we also have that  $B^4 = d(r-3)$  since Y is a variety of minimal degree. By Miyaoka's result (see [22]) we have that  $B^2 \cdot c_2 \ge 0$  and hence we have the inequality  $d \le \frac{24(r-1)}{r-3}$ .

(a) We now describe the cases when Y is a smooth variety of minimal degree. We have that  $r \ge 4$ .

*Case 1.* If r = 4, we have that  $Y = \mathbb{P}^4$ .

<u>*Case 2.*</u> If r = 5, we have that codimension of Y is one and degree is 2 which implies that Y is a smooth quadric hypersurface.

Suppose  $r \ge 6$ , we have that *Y* is a smooth rational normal scroll (which is abstractly a projective bundle over  $\mathbb{P}^1$ ) and is hence fibered over  $\mathbb{P}^1$ . Let this map from *Y* to  $\mathbb{P}^1$  be  $\phi$ . Composing this with  $\varphi$  we get a map  $\phi \circ \varphi : X \to \mathbb{P}^1$ . Therefore, *X* is fibered over  $\mathbb{P}^1$ . The general fibre is the inverse image of the general linear fiber of the smooth scroll and is hence irreducible by Bertini's theorem (see [10], Theorem 1.1). This along with generic smoothness implies that the general fibre of  $\phi$  is a

smooth threefold *G* with  $K_G = 0$  by adjunction. Let the general fibre of *Y* be denoted by *R* and that of *X* is denoted by *G*. We have the following exact sequence of cohomology of line bundles on *X*.

(2.16) 
$$0 \to H^0(B(-G)) \to H^0(B) \to H^0(B \otimes \mathscr{O}_G) \to H^1(B(-G)).$$

Notice that H - R is a base point free divisor in Y where H is a hyperplane section in Y. We have that Y is  $S(a_1, a_2, a_3, a_4)$  i.e, Y is the image of  $\mathbb{P}(\mathscr{E})$  where  $\mathscr{E}$  is the following vector bundle,

$$\mathbb{P}(\mathscr{E}) = \mathscr{O}(a_1) \oplus \mathscr{O}(a_2) \oplus \mathscr{O}(a_3) \oplus \mathscr{O}(a_4)$$

mapped to the projective space by  $|\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)|$ .

<u>*Case 3.*</u> For the cases r = 6 or r = 7 we use the fact that degree of  $\varphi$  is equal to the degree of  $\varphi|_G$  and then use Riemann-Roch theorem on the threefold *G* (see [15], Appendix A, Exercise 6.7) noticing the fact that  $K_G = 0$  and that  $B|_G$  is ample and base point free. This gives the upper bound  $6h^0(B|_G)$  since we have that  $B|_G \cdot c_2 \ge 0$  (see [22]). The lower bound 2 is due to the fact that *G* cannot be birational to  $\mathbb{P}^3$ .

Assuming *G* is regular and hence Calabi-Yau we have that  $h^0(B|_G) \ge \frac{1}{6}(B|_G)^3 + 1$  and hence we have  $d \le 6(h^0(B|_G) - 1)$ . The lower bound is obtained by Proposition 2.2, part (1) of [19].

<u>*Case 4.*</u> Suppose  $r \ge 8$ . Recall that H - R is base point free. We compute  $(H - R)^4 = H^4 - 4H^3R$ ;

(2.17) 
$$H^4 = \sum_{i=0}^3 a_i H^3 R \text{ and } r = \sum_{i=0}^3 a_i + 3.$$

So,  $r \ge 8$  gives  $\sum_{i=0}^{3} a_i \ge 5$  which gives  $(H - R)^4 > 0$  as *H* is ample. Hence, H - R is nef and big and consequently B - G is nef and big as well. Thus by Kawamata-Viehwag vanishing, we have that  $H^1(B - G) = 0$ . Hence  $\varphi|_G$  is given by the complete linear system  $|B|_G|$ .

Since G maps to  $F = \mathbb{P}^3$  we have that  $h^0(B|_G) = 4$ . Now, the degree of  $\varphi$  is also the degree of  $\varphi|_G$  for a general fibre G. Hence by a result of Gallego and Purnaprajna (see [11], Theorem 1.6) we have that  $2 \le d \le 18$ .

(b) Now we assume that *Y* is singular.

<u>*Case 1.*</u> As before, if r = 5 then Y is a singular quadric.

<u>*Case 2.*</u> Suppose the image Y of X under the morphism defined by |B| is a singular variety. Then by Theorem 1.8, Y is a cone over a smooth variety of minimal degree with vertex V. Moreover, codim(V, Y) = 2 by Lemma 1.11. Hence Y can be either a triple cone over a rational normal curve or a double cone over the Veronese surface in  $\mathbb{P}^5$ .

Suppose Y is a cone over a rational normal curve. Let R be a general linear subspace of Y and G its inverse image. By Bertini (see [10], Theorem 1.1), G is irreducible and  $d = B^3 \cdot G$ . Moreover, using the fact that the codimension of the singular locus of Y is exactly 2, we have  $B = deg(Y) \cdot G = (r-3)G$  by Lemma 1.9. Hence  $d \ge (r-3)^3$ . Using the previously proved upper bound we have,

$$(r-3)^3 \le 24(r-1)$$

and hence  $r \leq 8$ .

Now we prove Theorem A using the previous theorem. We notice that part (2) of the next theorem holds for both hyperkähler and Calabi-Yau four-folds in dimension four since it only requires a regular fourfold with trivial canonical bundle.

**Theorem 2.4.** *Let X be a four-fold with trivial canonical bundle and let A be an ample line bundle on X. then,* 

- (1) *nA* is very ample and embeds X as a projectively normal variety for  $n \ge 16$ .
- (2) If  $H^1(\mathcal{O}_X) = 0$  then nA is very ample and embeds X as a projectively normal variety for  $n \ge 15$ .

*Proof of (1).* By the result of Kawamata (see [18]), we have that on a fourfold with trivial canonical bundle if *A* is ample then *nA* is base point free for  $n \ge 5$ . Now using CM lemma (see Lemma 1.6) we can easily prove that *nA* satisfies the property  $N_0$  for  $n \ge 21$ .

If we set B = 5A then 20A = 4B and it satisfies the property  $N_0$  by Lemma 2.1. Using Lemma 2.2, Lemma 1.6 and Observation 1.3, we can see that the following map

$$H^0(nkA) \otimes H^0(nA) \longrightarrow H^0((nk+n)A)$$

is surjective for  $k \ge 2$  and  $16 \le n \le 19$ . So we are left to check the surjectivity of the multiplication map  $H^0(nA) \otimes H^0(nA) \longrightarrow H^0(2nA)$  for  $16 \le n \le 19$ . We just prove it for n = 16. The proof is similar for the other three cases.

For n = 16, the following maps surjects by Lemma 2.2 and Lemma 1.6,

~

(2.18) 
$$H^0(16A) \otimes H^0(5A) \to H^0(21A) \text{ and } H^0(21A) \otimes H^0(5A) \to H^0(26A).$$

Therefore, by Observation 1.3 we need to show that  $H^0(26A) \otimes H^0(6A) \longrightarrow H^0(32A)$  is surjective which follows from Lemma 1.6 as well.

*Proof of (2).* Suppose  $H^1(\mathcal{O}_X) = 0$ . We just need to show that 15A satisfies the property  $N_0$ .

Let B = 5A which is ample and base point free (see [18]). By Corollary 1.8, we know that 3B is projectively normal unless the image of the morphism induced by the complete linear series |B| is a variety of minimal degree. Thus, we aim to show that the image of the morphism induced by the complete linear series |5A| is not a variety of minimal degree. Applying Riemann-Roch we get,

(2.19) 
$$h^{0}(5A) = \frac{625}{24}A^{4} + \frac{25}{24}A^{2} \cdot c_{2} + 2 \ge 28.$$

Now suppose that the image is a variety of minimal degree. However, since the codimension of the image is  $\geq 24$ , by Theorem 2.3, we have that the image cannot be a quadric hypersurface or a cone over the veronese embedding of  $\mathbb{P}^2$  in  $\mathbb{P}^5$  or a cone over a rational normal curve. Hence the image is a smooth rational normal scroll.

Let  $h^0(B) = r + 1$ . Hence the degree of the image is r - 3. Also, let the degree of the finite morphism given by the complete linear series of *B* be *d*. We know by Theorem 2.3 that

$$d \le \frac{24(r-1)}{r-3}.$$

Using  $h^0(B) \ge 28$  we have that  $r \ge 27$  and hence  $d \le 26$ .

Since the image of the morphism is a smooth rational normal scroll of dimension 4, we can choose a general  $\mathbb{P}^3 = R$  and take the pullback of the divisor *R* under the morphism induced by the complete linear series |B| and call it *G*. The degree of the morphism restricted to *G* is again *d*. Since the degree of *R* in the image is 1 we have that  $d = B^3 \cdot G = 125A^2 \cdot G \ge 125$  (since *A* is ample and *G* is effective) contradicting  $d \le 26$ . Hence the image cannot be variety of minimal degree.  $\Box$ 

### MUKHERJEE AND RAYCHAUDHURY

#### 3. Proof of Theorem B

Let *X* be a hyperkähler variety of dimension 2*n* and let *B* be an ample and globally generated line bundle on *X*. In this section,  $\varphi_B$  will always denote the morphism induced by the complete linear series |B|. The aim is to study the projective normality of  $B^{\otimes 2n-1}$ . We do this by Theorem 1.8 that require us to analyze the case when  $\varphi_B$  maps *X* onto a variety of minimal degree. Recall that a variety of minimal degree is either (1) a quadric hypersurface, or (2) a smooth rational normal scroll, or (3) cone over a smooth rational normal scroll, or (4) cone over the Veronese embedding of  $\mathbb{P}^2$  inside  $\mathbb{P}^5$  ([6]). The following lemma eliminates a few cases.

**Lemma 3.1.** Let *B* be an ample and globally generated line bundle on a hyperkähler variety *X* of dimension 2*n*. Suppose the morphism  $\varphi_B$  induced by the complete linear series |B| maps *X* onto a variety of minimal degree *Y*. Then,

- (1) If Y is a quadric hypersurface then  $h^0(B) = 2n + 2$ .
- (2) Y can never be a smooth rational normal scroll.
- (3) If Y is a cone over a smooth rational normal scroll then the codimension of its singular locus is two i.e. Y is obtained by taking cones over a smooth rational normal curve.
- (4) If Y is a cone over the Veronese embedding of  $\mathbb{P}^2$  inside  $\mathbb{P}^5$  then  $h^0(B) = 2n + 4$ .

*Proof.* (1) and (4) are obvious. (3) comes from Lemma 1.11. We give the proof of (2) below.

To prove (2) we argue by contradiction. Suppose the image is a smooth rational normal scroll. Since a smooth scroll admits a morphism to  $\mathbb{P}^1$  we have a composed morphism from X to  $\mathbb{P}^1$ . Take the Stein factorization of this morphism which has connected fibres and notice that since X is smooth this further factors through a normalization. So we get a morphism from X to a normal base of dimension 1 (hence smooth in this case) with connected fibres which contradicts Matsushita's result on the fibre space structure of a holomorphic symplectic manifold (see Theorem 1.15).  $\Box$ 

We give two definitions below that we will use later in this note.

**Definition 3.2.** For a given hyperkähler variety X, we define the following two polynomials,

$$RR_X(x) = \sum_{i=0}^n \frac{a_i}{(2i)!} (x)^i$$
 and  $R_X(x) = RR_X(x) - \frac{a_n}{(2n)!} x^n$ 

where  $a_i = C(td_{2n-2i})$ . (Note: these polynomials depend only on the deformation type of X.)

**Definition 3.3.** For a given hyperkähler variety X with Beauville form  $q_X$ , we define the constant  $\alpha_X$  as below,

$$\alpha_X = \min \Big\{ q_X(A) \Big| A \text{ is an ample line bundle on } X \Big\}.$$

Our next task is to find an upper bound for  $deg(\varphi_B)$  for an ample and base point free line bundle *B*. The next result in fact just uses the fact that *B* is base point free and the induced morphism  $\varphi_B$  is generically finite.

**Lemma 3.4.** Let X be a hyperkähler manifold of dimension 2n. Assume  $R_X|_{\mathbb{Z}\geq 0}$  is increasing and  $R_X(\alpha_X) > 2n$ . Then, for any globally generated line bundle B on X, such that  $\varphi_B$  is generically finite,  $deg(\varphi_B) < (2n)!$ .

*Proof.* Let  $Y = Im(\varphi_B)$ . We have the equation  $B^{2n} = deg(\varphi_B) \cdot deg(Y)$  to begin with. Note that,

$$(3.1) \qquad \qquad \deg(Y) \ge 1 + \operatorname{codim}(Y) \implies \deg(Y) \ge h^{\circ}(B) - 2n$$

since  $codim(Y) = h^0(B) - 2n$ . Thus, we get,

$$(3.2) B^{2n} \ge deg(\varphi_B)(h^0(B) - 2n).$$

Recall that  $B^{2n} = c_X(q_X(B))^n$  and notice that  $h^0(B) = RR_X(q_X(B)) = R_X(q_X(B)) + \frac{a_n}{(2n)!}(q_X(B))^n$ . Since,  $a_n = c_X$ , using (3.2) we get,

$$c_X(q_X(B))^n \ge deg(\varphi_B) \left( R_X(q_X(B)) + \frac{a_n}{(2n)!} (q_X(B))^n - 2n \right)$$
  
$$\implies c_X(q_X(B))^n \left( 1 - \frac{deg(\varphi_B)}{(2n)!} \right) \ge deg(\varphi_B) (R_X(q_X(B)) - 2n) \ge deg(\varphi_B) (R_X(\alpha_X) - 2n).$$

That concludes the proof since the last term is strictly greater than zero by hypothesis and  $q_X(B) > 0$  since *B* is nef and big.

**Remark 3.5.** If all Todd classes of the hyperkähler variety X is fakely effective then  $R_X|_{\mathbb{Z}\geq 0}$  is increasing. In particular, it is satisfied for all known examples of hyperkähler varieties, except O'Grady's 10 dimensional example  $\mathcal{M}_{10}$  (see [4], Theorem 1.8) remaining unknown, which is also clear from their explicit Riemann-Roch expressions.

The remark above leads to the following consequence.

**Remark 3.6.** The hypothesis of Lemma 3.4 are satisfied for all known examples of hyperkähler varieties X of dimension  $2n \ge 4$  (see [4], Theorem 1.8), except O'Grady's 10 dimensional example  $\mathcal{M}_{10}$ .

*Proof.* Thanks to the previous remark, it is enough to show that  $R_X(\alpha_X) > 2n$ .

If X is either type  $K3^{[n]}$  or  $K^n(T)$  then  $\alpha_X \ge 2$ . Since  $R_X|_{\mathbb{Z}\ge 0}$  is increasing, we have,

$$(3.3) R_X(\alpha_X) \ge RR_X(2) - \frac{c_X 2^n}{(2n)!}.$$

If X is of type  $K3^{[n]}$  (resp.  $K^n(T)$ ), using the Riemann-Roch expression 1.3 (resp. 1.4) and  $n \ge 2$  we have the following,

(3.4) 
$$RR_X(2) - \frac{c_X 2^n}{(2n)!} = \binom{n+2}{n} - \frac{1}{n!} > 2n \quad \left(\text{resp. } RR_X(2) - \frac{c_X 2^n}{(2n)!} = (n+1)^2 - \frac{1}{n!} > 2n\right).$$

Same argument works for  $\mathcal{M}_6$  as well since its Riemann-Roch expression is same as that of  $K^3(T)$ . That concludes the proof of the assertion.

The upper bound for the degree has the following consequence on the secant lines of an embedding of a *K*3 surface. Even though it has nothing to do with our present purpose, it still might be of independent interest.

**Corollary 3.7.** Let *S* be a K3 surface and *B* be a very ample line bundle. Consider the projective embedding of *S* in  $\mathbb{P}^{h^0(B)-1}$  and the closed subvariety of  $Gr(2, h^0(B))$  consisting of lines that intersect the K3 surface at a subscheme of length at least 2. Then a general such line intersects *S* at a subscheme of length  $\leq 7$ .

*Proof.* Given the above conditions we construct a generically finite morphism f from  $X = S^{[2]}$  to  $Gr(2, h^0(B))$ . Given a point on X we take the length 2 subscheme it defines on S and send it to the linear span of the length two subscheme inside  $\mathbb{P}^{h^0(B)-1}$ . Since a general such line does not lie on S, it intersects S at finitely many points. So a general point in the image of the morphism f has got finite fibers. Hence f is a generically finite morphism.

Also this morphism is given by the complete linear series  $B - \delta$  where  $2\delta$  is the class of the divisor in  $S^{[2]}$  that parametrizes non-reduced subschemes of length 2 on the K3 surface S. By Lemma 3.4 and Remark 3.6, we have  $deg(f) \le 23$ . If a line intersects S at k points then the line has  $\binom{k}{2}$  preimages under the morphism f. Thus for a general line  $\binom{k}{2} \le 23$  and hence  $k \le 7$ .

In the next Lemma, we will use the upper bound for  $deg(\varphi_B)$  to put more restrictions on the image of  $\varphi_L$  when it maps onto a variety of minimal degree.

**Lemma 3.8.** Let X be a hyperkähler manifold of dimension  $2n \ge 4$  for which  $R_X|_{\mathbb{Z}\ge0}$  is increasing,  $R_X(\alpha_X) > 2n$  and  $RR_X(\alpha_X) \ge 4n$ . Then for any ample and globally generated line bundle B on X,  $\varphi_B$  can never map X onto a variety that is obtained by taking cones over a rational normal curve.

*Proof.* Suppose  $Y = Im(\varphi_B)$  is a variety of minimal degree that is obtained by taking cones over a rational normal curve and  $d = deg(\varphi_B)$ . Therefore Y is singular in codimension two. Let G be the inverse image of a general linear subspace of R of codimension 1 in Y. Notice that G is irreducible by Bertini's theorem.

We have  $B^{2n-1} \cdot G = d$ . Using Lemma 1.10, we deduce that *B* can be written as the pullback of  $deg(Y) \cdot R$ . Thus, *G* is ample and  $d = deg(Y)^{2n-1} \cdot G^{2n} \ge deg(Y)^{2n-1}$ .

Now we use the fact that  $deg(Y) = (RR_X(q_X(B)) - 2n)$  and d < (2n)! (see Lemma 3.4 and Remark 3.6) that leads us to the following inequality,

(3.5) 
$$(RR_X(q_X(B)) - 2n)^{2n-1} < (2n)!$$

which is absurd. Indeed, by our assumption,  $RR_X(q_X(B)) - 2n \ge RR_X(\alpha_X) - 2n \ge 2n$ .

Notice that the proof above also shows the following.

**Remark 3.9.** Let X be a hyperkähler manifold of dimension  $2n \ge 4$  for which  $R_X|_{\mathbb{Z}\ge0}$  is increasing and  $R_X(\alpha_X) > 2n$ . Let B be an ample and globally generated line bundle for which either  $h^0(B) \ge 4n$ , or  $B^{2n} < \alpha_X^n (h^0(B) - 2n)^{2n}$ . Then  $\varphi_B$  can never map X onto a variety that is obtained by taking cones over a smooth rational normal curve.

Combining Lemmas 3.1, 3.8 and Remarks 3.6 and 3.9 we get the following.

**Proposition 3.10.** Let X be a hyperkähler manifold of dimension  $2n \ge 4$  and let B be an ample and globally generated line bundle on X. If X is deformation equivalent to either  $K^n(T)$  or  $\mathcal{M}_6$  then the morphism  $\varphi_B$  given by the complete linear series |B| will never map X onto a variety of minimal degree. If X is of type  $K3^{[n]}$  and  $\varphi_B$  maps X onto a variety Y of minimal degree then either,

- (1) X is of type  $K3^{[2]}$ ,  $q_X(B) = 2$ ,  $deg(\varphi_B) = 6$  and Y is a quadric hypersurface (possibly singular) inside  $\mathbb{P}^5$  which can not be obtained by taking cones over any rational normal scroll, or
- (2) X is of type  $K3^{[3]}$ ,  $q_X(B) = 2$ ,  $deg(\varphi_B) = 30$  and Y is a variety embedded inside  $\mathbb{P}^9$  that is obtained by taking cones over the Veronese embedding of  $\mathbb{P}^2$  inside  $\mathbb{P}^5$ .

*Proof.* To start with, note that  $\alpha_X \ge 2$ . Suppose *X* is of type  $K^n(T)$  or  $\mathcal{M}_6$ . By Riemann-Roch, we get  $h^0(B) = RR_X(q_X(B)) \ge RR_X(2) > 2n + 4$ . Consequently, by Lemma 3.1,  $Im(\varphi_B)$  can only be a variety that is obtained by taking cones over a rational normal curve. But that is impossible by Lemma 3.8 since  $RR_X(2) > 4n$ .

Now, assume X is of type  $K3^{[n]}$ . We can argue exactly like the previous paragraph to conclude that  $\varphi_B$  will never map X onto a variety of minimal degree if  $n \ge 5$ .

We deal with the case n = 2, 3 and 4 separately and we will use Lemma 3.1. Note that  $q_X(B)$  is even, say  $q_X(B) = 2k$  for some positive integer k.

Suppose n = 2. Note that  $h^0(B) > 8$  if  $q(B) \ge 4$  and  $B^{2n} < 2^2(h^0(B) - 4)^4$  if q(B) = 2. Consequently by Remark 3.9, Y can not be obtained by taking cones over rational normal curve. The equation  $RR_X(2k) = 2n + 2$  has only one positive even integer solution k = 1 in which case  $q_X(B) = 2$ ,  $deg(\varphi_B) = 6$  and Y is a quadric hypersurface in  $\mathbb{P}^5$ .  $RR_X(2k) = 2n + 4$  has no integer solution.

Suppose n = 3. Argument similar to that of the case n = 2 yields that Y can not be obtained by taking cones over rational normal curve. Moreover,  $RR_X(2k) = 2n + 2$  has no solution and  $RR_X(2k) = 2n+4$  has only one positive integer solution k = 1 in which case  $q_X(B) = 2$ ,  $deg(\varphi_B) = 30$ and Y is a variety of minimal degree in  $\mathbb{P}^9$  obtained by taking veronese embedding of  $\mathbb{P}^2$  inside  $\mathbb{P}^5$ .

Similar argument shows that  $\varphi_B$  can not map  $K3^{[4]}$  onto a variety of minimal degree.  $\Box$ 

Now we are ready to give the proof of Theorem B.

**Theorem 3.11.** Let X be a projective hyperkähler variety of dimension  $2n \ge 4$  that is deformation equivalent to  $K3^{[n]}$ ,  $K^n(T)$  or  $\mathcal{M}_6$ . Let B be an ample and globally generated line bundle on X. Then the following happens;

- (1)  $B^{\otimes l}$  is projectively normal for  $l \geq 2n$ .
- (2)  $B^{\otimes 2n-1}$  is projectively normal unless:
  - (a) n = 2,  $X = K3^{[2]}$  and  $\varphi_B$  maps X onto a quadric (possibly singular) inside  $\mathbb{P}^5$ . In this case  $q_X(B) = 2$ ,  $deg(\varphi_B) = 6$ , or
  - (b) n = 3,  $X = K3^{[3]}$  and  $\varphi_B$  maps X onto a variety of minimal degree inside  $\mathbb{P}^9$  which is obtained by taking cones over the Veronese embedding of  $\mathbb{P}^5$  inside  $\mathbb{P}^5$ . In this case  $q_X(B) = 2$ ,  $deg(\varphi_B) = 30$ .

Hence if X is as above and B does not satisfy cases 2(a) or 2(b) then a general curve section of |B| is non-hyperelliptic.

*Proof.* To prove (1) we simply notice that  $h^0(B) \ge 2n + 2$  by the Riemann-Roch formula on *X*. The assertion follows by Lemma 2.1. (2) follows directly by Corollary 1.8, and Proposition 3.10. The statement on non-hyperellipticity of a general curve section *C* follows from the fact that  $B^{d-1}|_C = K_C$  by adjunction and that a very ample line bundle restricts to a very ample line bundle on a closed immersion.

We finish by the following example of Debarre (see [5]). It shows the existence of an ample and globally generated line bundle on a hyperkähler variety of  $K3^{[2]}$  type that induces a 6-1 map onto a variety of minimal degree by its complete linear series.

**Example 3.12** Let (S, L) be a polarized K3 surface with  $Pic(S) = \mathbb{Z}L$  and  $L^2 = 4$ . Then L is very ample and consequently we get a morphism  $\phi : S^{[2]} \to Gr(2, 4)$  to the Grassmannian.

Now, *L* induces a line bundle  $L_2$  on  $S^{[2]}$  and it is known that  $Pic(S^{[2]}) = \mathbb{Z}L_2 \oplus \mathbb{Z}\delta$ . Moreover, the pullback of the Plücker line bundle on the Grassmannian has class  $L_2 - \delta$  on  $S^{[2]}$ . Therefore, if (S, L) is general then it contains no line and consequently  $\phi$  will be finite of degree  $\binom{4}{2} = 6$ .  $\Box$ 

### References

- [1] Beauville, Arnaud. Variétés Kähleriennes dont la première classe de Chern est nulle. (French) J. Differential Geom. 18 (1983), no. 4, 755–782 (1984).
- Bogomolov, F. A. The decomposition of Kahler manifolds with a trivial canonical class. (Russian) Mat. Sb. (N.S.) 93(135) (1974), 573–575, 630.
- [3] Britze, M.; Nieper, M. A. *Hirzebruch-Riemann-Roch formulae on irreducible symplectic Kähler manifolds*. arXiv:math/0101062
- [4] Cao, Yalong; Jiang, Chen. Remarks on Kawamata's effective non-vanishing conjecture for manifolds with trivial first Chern classes. arXiv:1612.00184v2
- [5] Debarre, Olivier. Hyperkähler Manifolds. arXiv:1810.02087v1
- [6] Eisenbud, David; Harris, Joe. On varieties of minimal degree (a centennial account). Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), 3–13, Proc. Sympos. Pure Math., 46, Part 1, Amer. Math. Soc., Providence, RI, 1987.
- [7] Ellingsrud, Geir; Göttsche, Lothar; Lehn, Manfred. On the cobordism class of the Hilbert scheme of a surface.
  J. Algebraic Geom. 10 (2001), no. 1, 81–100.
- [8] Ferraro, Rita. Weil divisors on rational normal scrolls. Geometric and combinatorial aspects of commutative algebra (Messina, 1999), 183–197, Lecture Notes in Pure and Appl. Math., 217, Dekker, New York, 2001.
- [9] Fujiki, Akira. On the de Rham cohomology group of a compact Kähler symplectic manifold. Algebraic geometry, Sendai, 1985, 105–165, Adv. Stud. Pure Math., 10, North-Holland, Amsterdam, 1987.
- [10] Fulton, William and Lazarsfeld, Robert. Connectivity and its applications in algebraic geometry. Algebraic geometry (Chicago, Ill., 1980), pp. 26–92, Lecture Notes in Math., 862, Springer, Berlin-New York, 1981.
- [11] Gallego, Francisco Javier; Purnaprajna, B. P. Very ampleness and higher syzygies for Calabi-Yau threefolds. Math. Ann. 312 (1998), no. 1, 133–149.
- [12] Green, Mark L. The Canonical Ring of a variety of general type. Duke Math. J. 49 (1982), no. 4, 1087–1113.
- [13] Green, Mark L. Koszul cohomology and the geometry of projective varieties. J. Differential Geom. 19 (1984), no. 1, 125–171.
- [14] Gross, M.; Huybrechts, D.; Joyce, D. Calabi Yau manifolds and related geometries. Lectures from the Summer School held in Nordfjordeid, June 2001. Universitext. Springer-Verlag, Berlin, 2003. viii+239 pp. ISBN: 3-540-44059-3
- [15] Hartshorne, Robin. Algebraic geometry. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977.
- [16] Horikawa, Eiji. Algebraic Surfaces of General Type With Small c<sub>1</sub><sup>2</sup>. I Ann. of Math. (2) 104 (1976), no. 2, 357–387.
- [17] Huybrechts, Daniel. Compact hyper-K ahler manifolds: basic results. Invent. Math. 135 (1999), no. 1, 63–113.
- [18] Kawamata, Yujiro. On Fujitas freeness conjecture for 3-folds and 4-folds. Math. Ann. 308 (1997), no. 3, 491– 505.
- [19] Kanazawa, Atsushi; Wilson, P. M. H. Trilinear forms and Chern classes of Calabi-Yau threefolds. Osaka J. Math. 51 (2014), no. 1, 203–213.
- [20] Lazarsfeld, Robert. *Positivity in algebraic geometry. II.* Positivity for vector bundles, and multiplier ideals. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 49. Springer-Verlag, Berlin, 2004.
- [21] Matsushita, Daisuke. On fibre space structures of a projective irreducible symplectic manifold. Topology 38 (1999), no. 1, 79–83.
- [22] Miyaoka, Yoichi. The Chern classes and Kodaira dimension of a minimal variety. Algebraic geometry, Sendai, 1985, 449–476, Adv. Stud. Pure Math., 10, North-Holland, Amsterdam, 1987.

- [23] Mukherjee, Jayan; Raychaudhury, Debaditya. On the projective normality and normal presentation on higher dimensional varieties with nef canonical bundle. J. Algebra 540 (2019), 121–155.
- [24] Mumford, David. Varieties defined by quadratic equations. 1970 Questions on Algebraic Varieties (C.I.M.E., III Ciclo, Varenna, 1969) pp. 29–100 Edizioni Cremonese, Rome
- [25] Nieper, Marc A. Hirzebruch-Riemann-Roch formulae on irreducible symplectic Kähler manifolds. J. Algebraic Geom. 12 (2003), no. 4, 715–739.
- [26] Niu, Wenbo. On syzygies of Calabi-Yau varieties and varieties of general type. Adv. Math. 343 (2019), 756–788.
- [27] O'Grady, Kieran G. Desingularized moduli spaces of sheaves on a K3. J. Reine Angew. Math. 512 (1999), 49–117.
- [28] O'Grady, Kieran G. A new six-dimensional irreducible symplectic variety. J. Algebraic Geom. 12 (2003), no. 3, 435–505.
- [29] Oguiso, Keiji; Peternell, Thomas. *On polarized canonical Calabi-Yau threefolds*. Math. Ann. 301 (1995), no. 2, 237–248.
- [30] Saint-Donat, B. On projective models of K3 surfaces Amer. J. Math. 96 (1974), 602–639.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, LAWRENCE, KS 66045

E-mail address: j899m889@ku.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, LAWRENCE, KS 66045

E-mail address: debaditya@ku.edu