

MYCOTAXON

<http://dx.doi.org/10.5248/127.213>

Volume 127, pp. 213–219

January–March 2014

Three new species of *Pleurophragmium* from Yunnan, China

LI-GUO MA, JI-WEN XIA, YING-RUI MA, & XIU-GUO ZHANG*

Department of Plant Pathology, Shandong Agricultural University, Taian, 271018, China

*CORRESPONDENCE TO: zhxg@sdau.edu.cn, sdau613@163.com

ABSTRACT — Three new species are described and illustrated: *Pleurophragmium ellipsoideum* from dead branches of *Bauhinia acuminata*, *P. yunnanense* from dead branches of *Machilus salicina*, and *P. clavatum* from dead branches of *Beilschmiedia percoriacea*. All three species were collected from tropical forests in Yunnan Province, China.

KEY WORDS — anamorphic fungi, taxonomy

Introduction

Costantin (1888) established *Pleurophragmium* for a single species, *P. bicolor* Costantin. The genus is characterized by well defined single unbranched conidiophores with sympodial, terminal and intercalary, integrated, polyblastic, denticulate conidiogenous cells with pointed denticles and solitary apical and lateral conidia with pointed bases (Hughes 1958; Ellis 1971, 1976). These characters separate *Pleurophragmium* from similar genera — *Cordana* Preuss, *Cacumisporium* Preuss, *Spiropes* Cif., *Pseudospiropes* M.B. Ellis, *Minimelanolocus* R.F. Castañeda & Heredia, and *Dactylaria* Sacc. More than 20 species have been accepted in *Pleurophragmium*.

Several mycological papers dealing with many new species from China have been published recently (Zhang et al. 2009a,b, 2011, 2012; Liang et al. 2010, 2011; Dai 2011, 2012; Ma et al. 2011a,b, 2012). We collected three undescribed species of *Pleurophragmium* in our study on conidial fungi on dead wood from tropical forests in Yunnan Province, China. The specimens are deposited in HSAUP (Herbarium of the Department of Plant Pathology, Shandong Agricultural University) and HMAS (Mycological Herbarium, Institute of Microbiology, Chinese Academy of Sciences).

Taxonomy

Pleurophragmium ellipsoideum L.G. Ma & X.G. Zhang, sp. nov.

FIG. 1

MYCOBANK MB808631

Differs from *Pleurophragmium miniumbonatum* by its shorter and wider pale brown conidia with rounded apices, from *P. acutum* by its larger pale brown conidia with more septa, and from *P. obcampanuloides* by its wider conidia with more septa.

TYPE: China, Yunnan Province: the Forbidden Forest of Banna, on dead branches of *Bauhinia acuminata* L. (*Caesalpiniaceae*), 17 Oct. 2008, L.G. Ma (holotype HSAUP H0042; isotype HMAS 243411).

ETYMOLOGY: in reference to the ellipsoid conidial shape.



FIG. 1. *Pleurophragmium ellipsoideum* (holotype, HSAUPH0042).

a. Conidiophores and conidia. b. Conidiogenous cells with pointed denticles. c, d. Conidia.

Colonies on natural substratum effuse, brown. Mycelium partly immersed and partly superficial, composed of septate, branched, smooth, subhyaline to pale brown hyphae. Conidiophores well defined, single, unbranched, erect, straight or slightly flexuous, cylindrical, brown at the base, subhyaline to pale brown towards the apex, smooth, septate, thick-walled, 150–260 µm long, 2.5–3.5 µm wide at the apex, 3.5–6 µm wide at the base. Conidiogenous cells polyblastic, terminal and intercalary, integrated, sympodial, cylindrical, denticulate, smooth, thick-walled, subhyaline to pale brown, with numerous sharply pointed denticles towards the apex. Conidia solitary, apical and lateral, ellipsoid to obovoid, thick-walled, rounded at the apex, tapered to a point at the base, pale brown, smooth, 3-septate, 10–17 × 6.0–8.5 µm.

NOTES: *Pleurophragmium ellipsoideum* is similar to *P. miniumbonatum* (R.F. Castañeda et al.) R.F. Castañeda, *P. acutum* (Grove) M.B. Ellis, and *P. obcampanuloides* Matsush. in conidial shape. However, *P. miniumbonatum* can be easily distinguished from *P. ellipsoideum* by its longer and narrower versicolored conidia with umbonate apices (16–19 × 6–7 µm; Heredia-Abarca et al. 2007). *Pleurophragmium acutum* differs by its smaller hyaline 0-septate conidia (6–12 × 2–3 µm; Ellis 1976), while the conidia of *P. obcampanuloides* are 2-septate and narrower (10–18 × 5–7 µm; Matsushima 1995).

***Pleurophragmium yunnanense* L.G. Ma & X.G. Zhang, sp. nov.**

FIG. 2

MYCOBANK MB808632

Differs from *Pleurophragmium hippotrichoides* by its smooth conidia with more septa and from *P. ellipsoideum* by its broadly fusiform conidia with narrowed ends.

TYPE: China, Yunnan Province: the Forbidden Forest of Banna, on dead branches of *Machilus salicina* Hance (*Lauraceae*), 18 Oct. 2008, L.G. Ma (holotype HSAUP H0086; isotype HMAS 243412).

ETYMOLOGY: in reference to the province where the taxon was found.

Colonies on natural substratum effuse, brown. Mycelium partly immersed and partly superficial, composed of septate, branched, smooth, subhyaline to pale brown hyphae. Conidiophores well defined, single, unbranched, erect, straight or slightly flexuous, cylindrical, brown at the base, subhyaline to pale brown towards the apex, smooth, septate, thick-walled, 185–270 µm long, 2.5–3.3 µm wide at the apex, 3.5–4.5 µm wide at the base. Conidiogenous cells polyblastic, terminal and intercalary, integrated, sympodial, cylindrical, denticulate, smooth, thick-walled, subhyaline to pale brown, with numerous sharply pointed denticles towards the apex. Conidia solitary, apical and lateral, broadly fusiform, thick-walled, narrowed at the ends, tapered to a point at the base, pale brown, smooth, (2)–3-septate, 10–15 × 6–7 µm.

NOTES: *Pleurophragmium yunnanense* resembles *P. hippotrichoides* (Corda) M.B. Ellis in conidial shape, but *P. hippotrichoides* differs by its 0-septate



FIG. 2. *Pleurophragmium yunnanense* (holotype, HSAUPH0086).

a. Conidiophores and conidia. b. Conidiogenous cells with pointed denticles. c, d. Conidia.

verrucose conidia (Ellis 1976). *Pleurophragmium ellipsoideum* differs from *P. yunnanense* by its ellipsoid to obovoid conidia with rounded apices (see above).

***Pleurophragmium clavatum* L.G. Ma & X.G. Zhang, sp. nov.**

FIG. 3

MYCOBANK MB808633

Differs from *Pleurophragmium tritici* by its longer yellowish brown 0-septate conidia and from *P. acutum* by its larger yellowish brown verrucose conidia.

TYPE: China, Yunnan Province: the Forbidden Forest of Banna, on dead branches of *Beilschmiedia percoriacea* C.K. Allen (*Lauraceae*), 31 Oct. 2011, L.G. Ma (**holotype** HSAUPH2090; **isotype** HMAS 243416).

ETYMOLOGY: in reference to the clavate conidial shape.

Colonies on natural substratum effuse, brown. Mycelium partly immersed and partly superficial, composed of septate, branched, smooth, subhyaline to pale brown hyphae. Conidiophores well defined, single, unbranched, erect, straight or slightly flexuous, cylindrical, brown at the base, subhyaline to pale brown




FIG. 3. *Pleurophragmium clavatum* (holotype, HSAUPH2090).
a. Conidiophores and conidiogenous cells. b. Conidia.

towards the apex, smooth, septate, thick-walled, 75–160 µm long, 3.3–4 µm wide. Conidiogenous cells polyblastic, terminal and intercalary, integrated, sympodial, cylindrical, denticulate, smooth, thick-walled, subhyaline to pale brown, with numerous sharply pointed denticles towards the apex. Conidia solitary, apical and lateral, clavate to fusiform, thick-walled, narrowed at the ends, tapered to a point at the base, yellowish brown, verrucose, 0-septate, $9.5\text{--}16.5 \times 3.5\text{--}4.5$ µm.

NOTES: *Pleurophragmium clavatum* is similar to *P. tritici* M.B. Ellis and *P. acutum* in conidial shape. However, *P. acutum* can be separated by its smaller hyaline smooth conidia ($6\text{--}12 \times 2\text{--}3$ µm; Ellis 1976) and *P. tritici* by its shorter subhyaline or straw-coloured 0–1-septate conidia ($6\text{--}11 \times 3\text{--}4$ µm; Ellis 1976).

Acknowledgments

The authors are grateful to Dr. De-Wei Li and Dr. Bryce Kendrick for serving as pre-submission reviewers and for their valuable comments and suggestions. This project was supported by the National Natural Science Foundation of China (Nos. 31093440, 31230001) and the Ministry of Science and Technology of the People's Republic of China (No. 2006FY120100).

Literature cited

- Costantin J. 1888. Les mucédinées simples. Librairie Paul Klincksieck, Paris, France.
- Dai YC. 2011. A revised checklist of corticioid and hydroid fungi in China for 2010. *Mycoscience* 52: 69–79. <http://dx.doi.org/10.1007/s10267-010-0068-1>
- Dai YC. 2012. Polypore diversity in China with an annotated checklist of Chinese polypores. *Mycoscience* 53: 49–80. <http://dx.doi.org/10.1007/s10267-011-0134-3>
- Ellis MB. 1971. Dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey, England.
- Ellis MB. 1976. More dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey, England.
- Heredia-Abarca G, Castañeda-Ruiz RF, Arias RM, Saikawa M, Stadler M. 2007. Anamorphic fungi from submerged plant material: *Acumisporea verruculosa*, *Pleurophragmium aquaticum*, and *P. miniumbonatum*. *Mycotaxon* 101: 89–97.
- Hughes SJ. 1958. Revisiones hyphomycetum aliquot cum appendice de nominibus rejiciendis. *Can. J. Bot.* 36: 727–836. <http://dx.doi.org/10.1139/b58-067>
- Liang JF, Yang ZL, Xu J, Ge ZW. 2010. Two new unusual *Leucoagaricus* species (*Agaricaceae*) from tropical China with blue-green staining reactions. *Mycologia* 102: 1141–1152. <http://dx.doi.org/10.3852/09-021>
- Liang JF, Yang ZL, Xu DP. 2011. A new species in *Lepiota* (*Agaricaceae*) from China. *Mycologia* 103: 820–830. <http://dx.doi.org/10.3852/10-216>
- Ma J, Wang Y, O'Neill NR, Zhang XG. 2011a. A revision of the genus *Lomaanthes*, with the description of a new species. *Mycologia* 103: 407–410. <http://dx.doi.org/10.3852/10-176>
- Ma J, Wang Y, Ma LG, Zhang YD, Castañeda-Ruiz RF, Zhang XG. 2011b. Three new species of *Neosporidesmium* from Hainan, China. *Micol. Prog.* 10: 157–162. <http://dx.doi.org/10.1007/s11557-010-0685-2>

- Ma LG, Ma J, Zhang YD, Zhang XG. 2012. *Spadicoides camelliae* and *Diplococcium livistonae*, two new hyphomycetes on dead branches from Fujian Province, China. *Mycoscience* 53: 25–30. <http://dx.doi.org/10.1007/s10267-011-0138-z>
- Matsushima T. 1995. Matsushima mycological memoirs 8. Published by the author, Kobe, Japan.
- Zhang K, Ma LG, Zhang XG. 2009a. New species and records of *Shrungabeeja* from southern China. *Mycologia* 101: 573–578. <http://dx.doi.org/10.3852/09-006>
- Zhang K, Ma J, Wang Y, Zhang XG. 2009b. Three new species of *Piricaudiopsis* from southern China. *Mycologia* 101: 417–422. <http://dx.doi.org/10.3852/08-147>
- Zhang YD, Ma J, Wang Y, Ma LG, Castañeda-Ruiz RF, Zhang XG. 2011. New species and record of *Pseudoacrodictys* from southern China. *Micol. Prog.* 10: 261–265. <http://dx.doi.org/10.1007/s11557-010-0696-z>
- Zhang YD, Ma J, Ma LG, Zhang XG. 2012. Two new species of *Taeniolina* from southern China. *Micol. Prog.* 11: 71–74. <http://dx.doi.org/10.1007/s11557-010-0729-7>