

Conidial fungi from the semi-arid Caatinga biome of Brazil. New species and new records of *Helicosporium*

ALISSON CARDOSO RODRIGUES DA CRUZ
alissonbio@yahoo.com.br

TASCIANO DOS SANTOS SANTA IZABEL
tazuefsbot@gmail.com

SHEILA MIRANDA LEÃO-FERREIRA
sheila1leao@yahoo.com.br

&

LUÍS FERNANDO PASCHOLATI GUSMÃO
lgusmao@uefs.br

*Departamento de Ciências Biológicas, Laboratório de Micologia
Universidade Estadual de Feira de Santana*

Av. Transnordestina, S/N – Novo Horizonte, 44036-900. Feira de Santana, BA, Brazil

Abstract —A new helicosporous fungus, *Helicosporium vesiculiferum*, is described based on morphological characters. It was found on a decaying twig of an unidentified dicotyledonous plant from the semi-arid region, Northeast of Brazil. The new species is illustrated with light micrographs and scanning electron microscopy. *Helicosporium vesiculiferum* is characterized by the presence of vesicles on apex of the conidiophores. Seven helicosporous species are briefly described, of which *H. virescens* and *H. aureum* are new records from Brazil, *H. panacheum* and *H. gracile* are reported for the first time in South America, and *H. nizamabadense* is new for the American continent. A key to all *Helicosporium* species is included.

Key words — anamorphic fungi, taxonomy

Introduction

Helicosporous fungi are defined by their coiled asexual spores (Tsui & Berbee 2006) and are considered aero-aquatic (Dix & Webster 1995). They are commonly found on substrates with abundant water (Mercado-Sierra 1982). In the present study these fungi were found in a semi-arid region, demonstrating the plasticity and survival capacity of helicosporous fungi in an inhospitable environment with low water availability and high temperatures.

Since Link (1809) described the first helicosporous species, *Helicomycetes roseus*, several studies have been devoted to this group (e.g., Morgan 1892;

Linder 1929; Moore 1953, 1955; Goos 1980, 1985a,b,c, 1986, 1989, 1990). There are about 240 described species in 45 genera (Saccardo 1886, Linder 1929, Goos 1987, Zhao et al. 2007).

Phylogenetic studies are clarifying the relationship between the major genera of helicosporous fungi and their teleomorph connections. Tsui et al. (2006) analyzed rDNA of species of *Helicomycetes* Link, *Helicoma* Corda, and *Helicosporium* Nees and verified that although all were related to *Tubeufia* Penz. & Sacc. sensu M.E. Barr, they are not monophyletic. Tsui & Berbee (2006) confirmed and extended the polyphyletism to *Helicodendron* Peyronel and *Helicoon* Morgan and suggested that the convergent form of the conidia is a possible adaptation to dispersion in aquatic environments.

Helicosporium is characterized by the presence of long and conspicuous conidiophores and conidia that are relatively thin-walled and hygroscopic (Goos 1989). The species are mainly distinguished by the morphology of conidia (diameter, filament width and number of coils), conidiophores, conidiogenous cells, and colony color (Goos 1989). Currently 21 species are included in the genus (Zhao et al. 2007)

Up to now, nine helicosporous fungi have been reported for Brazil: *Dichotomophthoropsis nympheae* (F.V. Rand) M.B. Ellis (Cavalcanti & Milanez 2007), *Hiospira jambosae* Bat. et al. (Batista et al. 1964), *Helicoma bambusae* Henn. (Hennings 1902), *H. palmarum* G.Z. Zhao et al. (Samuels et al. 1978, as *Helicomycetes* sp.), *Helicomycetes roseus* Link (Goos 1985b), *Helicosporium griseum* and *H. guianense* Linder (CBS 2009), *H. pannosum* (Sivanesan 1984), and *Xenosporium berkeleyi* (M.A. Curtis) Piroz. (Mendes et al. 1998, as *Xenosporella berkeleyi* (M.A. Curtis) Linder).

Previous studies of leaf litter conidial fungi from Caatinga biome have demonstrated the richness of this region (Castañeda-Ruiz et al. 2006; Cruz et al. 2007 a, b; Gusmão et al. 2008; Leão-Ferreira et al. 2008). During a recent survey an interesting specimen of *Helicosporium* was found. It does not resemble hitherto described species and is proposed here as new.

Materials and methods

The Program of Research in Biodiversity of the Brazilian semi-arid (PPBIO/Semi-árido) is investigating the biodiversity in Caatinga biome. This biome is considered part of a phytogeographical unit of South America called the neotropical seasonally dry tropical forests (Pennington et al. 2000, Prado 2000). It is characterized by high annual average temperatures (26–28°C) and a low precipitation (300–1000 mm/year) that is usually concentrated within 3–5 months; droughts are common (Sampaio 1995).

Samples of plant litter were collected in separate paper bags and taken to the laboratory where each was incubated at 25°C in Petri dishes within 170 L plastic chambers containing 200 ml sterile water and 2 ml glycerol. The plant material was screened at regular intervals for microfungi. Mounts were prepared in polyvinyl alcohol-glycerol

(8 g per 100 ml H₂O, 5 ml glycerol) and deposited in the Herbarium of Universidade Estadual de Feira de Santana (HUEFS).

Taxonomy

Helicosporium vesiculiferum A.C. Cruz & Gusmão sp. nov.

FIGS. 1–9

MYCOBANK MB512999

COLONIAE in substrato naturali effusae, albae. MYCELIUM partim superficiale et partim in substrato immersum, ex hyphis ramosis, septatis, pallide brunneis laevibus, 3–6 µm latis compositum. CONIDIOPHORA ex hyphis oriunda, macronematosae, mononematosae, 4–12-septatae, simplicia vel interdum ramosa prope basem, erecta, recta vel flexuosa, laevia, brunnea, pallide brunnea vel subhyalina ad apicem, 32.5–155 × 3–7.5 µm; vesicula in apice, plerumque praesentia, globosa, subhyalina, 5–9 × 4.5–8 µm; rami 1–2-septatae, laeves, brunnea vel pallide brunnea ad apicem, 10–35(–80) × 2.5–6 µm. CELLULAE CONIDIOGENAE polyblasticae, in conidiophoris incorporatae, intercalares et terminales, cylindricae, laevia, brunneae vel pallidae brunneae; denticulae 0.8–2 × 0.7–2 µm. CONIDIA holoblastica, solitaria, sicca, acropleurogena, 10–15-septata, helicoidea, laevia, hyalina, 11–18 µm diam, filamenta conidica 1–1.5 µm crassa, in 2–3.5 spiris convolute.


HOLOTYPE: BRAZIL. BAHIA: Senhor do Bonfim, in ramulis emortuis angiospermae ignotae, 25.IX.2006, A.C.R. Cruz, HUEFS 129360.

ETYMOLOGY: Latin, *vesiculiferum* = carrying vesicles.

COLONIES on natural substrate effuse, white. MYCELIUM superficial and immersed. HYPHAE septate, branched, smooth, pale brown, 3–6 µm diam. CONIDIOPHORES distinct, mononematous, 4–12-septate, simple or occasionally branched near the base, erect, straight or flexuose, smooth, brown, becoming pale brown towards subhyaline apex, 32.5–155 × 3–7.5 µm; vesicle usually present at the apex, globose, pale brown to subhyaline, 5–9 × 4.5–8 µm; branches 1–2-septate, smooth, brown to pale brown, 10–35(–80) × 2.5–6 µm. CONIDIOGENOUS CELLS polyblastic, arise from creeping mycelium, from conidiophores or branches, intercalary or terminal, integrated, cylindrical, smooth, brown to pale brown; denticles, 0.8–2 × 0.7–2 µm. CONIDIA holoblastic, acropleurogenous, 10–15-septate, helical, smooth, hyaline, 11–18 µm diam., conidial filament 1–1.5 µm wide, coiled 2–3.5 times.

OTHER MATERIAL EXAMINED: BRAZIL. BAHIA: Morro do Chápeu, on decaying twig, 12.II.2009, T.S. Santa Izabel, HUEFS 141556.

COMMENTS: *Helicosporium vesiculiferum* is distinguished from other species in the genus by the presence of a vesicle at the conidiophore apex. Only four species produce conidia acropleurogenously: *H. gracile*, *H. griseum*, *H. guianense*, and *H. lumbricopsis*. *Helicosporium griseum* and *H. lumbricopsis* have conidiophores that are branched and often anastomosing, and can be differentiated easily from *H. vesiculiferum* (Berkeley 1874, Linder 1929). *Helicosporium guianense* possess branches along the tall conidiophores (Linder 1929). *Helicosporium gracile* is distinguished by yellow colony and absence of vesicles (Morgan 1892, Goos 1989).



FIGS. 1–6. Light micrographs of *Helicosporium vesiculiferum* (from holotype).
 1. General aspect on natural substrate; 2–3. Conidiophores with and without vesicles;
 4–5. Detail of the conidiophore apex; 6. Conidium
 (Bars 1–3= 20 μ m, 4–6= 5 μ m)

Other species found in semi-arid region

Helicosporium aureum (Corda) Linder, Ann. Mo. bot. Gdn. 16: 279. 1929.

TEL.: *Acanthostigma scopulum* (Cooke & Peck) Peck, Bull. New York State Mus. 1: 22. 1887.

COLONY yellow. CONIDIOPHORES 147–255 $\times$ 4–5.5 μ m. CONIDIOGENOUS CELLS tooth-like or bladder-like, 4.5–13.5 $\times$ 2.5–3.5 μ m. CONIDIA pleurogenous,




FIG. 7–9. Scanning Electron Microscope micrographs of *Helicosporium vesifuliferum*.

7. Conidiophore; 8. Vesicle; 9. Detail of the secession of conidium.

(Bars fig. 7= 10 μm , 8= 3 μm , 9= 2 μm)

9–20-septate, 10.5–15 μm diam., filament width 1–1.7 μm , number of coils 2.5–4 times.

SPECIMENS EXAMINED: BRAZIL. BAHIA: Pilão Arcado, on dead bark, 16.II.2007. A.C.R. Cruz, HUEFS 129346; Senhor do Bonfim, “Carrapichel”, on decaying twig, 9.X.2006, A.C.R. Cruz, HUEFS 129347.

DISTRIBUTION: Argentina, Australia, Austria, Belize, Brazil, China, Czech Republic, formerly USSR, Japan, Pakistan, Panama, New Zealand, South Africa, USA (CBS 2009, Goos 1989, Farr & Rossman 2009, Morris 1978, Romero & Pildain 2004, Zhao et al. 2007).

Helicosporium aureum is similar to *H. decumbens*, *H. guianense* and *H. hiospiroides* B.S. Reddy et al. in possessing discrete, polyblastic conidio-

genous cells. *Helicosporium decumbens* and *H. hiospiroides* are distinguished by conidia diameter. *Helicosporium guianense* develops branches along its conidiophores. The Brazilian specimens are consistent with the descriptions given by Linder (1929), Goos (1989), and Zhao et al. (2007) regarding the absence of apical branches on conidiophores. *Helicosporium aureum* is recorded for the first time for Brazil.

Helicosporium gracile (Morgan) Linder, Ann. Missouri Bot. Gard. 16: 281. 1929.

COLONY yellow. CONIDIOPHORES 25.5–150 × 3–6 µm. CONIDIOGENOUS CELLS tooth-like. CONIDIA acropleurogenous, 12–15-septate, 12–15µm diam., filament width 1–1.5 µm, number of coils 2–3.5 times.

SPECIMENS EXAMINED: BRAZIL. BAHIA: Morro do Chapéu, on dead leaves, 07.X.2008. T.S. Santa Isabel, HUEFS 141555.

DISTRIBUTION: Africa, Brazil, China, USA (Goos 1989, Linder 1929, Zhao et al. 2007).

The characteristics observed on the examined specimen are included within the morphological variation for the species (Linder 1929, Goos 1989, Zhao et al. 2007). *Helicosporium gracile* is closely related to *H. guianense*, *H. aureum*, and *H. virescens*. However, *H. guianense* differs from *H. gracile* by consistent branches along the conidiophores. *Helicosporium aureum* can be separated by bladder-like conidiogenous cells and larger conidiophores and *H. virescens* has tooth or bladder-like conidiogenous cells and larger conidiophores (Goos 1989, Zhao et al. 2007). This species is reported for the first time in South America.

Helicosporium griseum Berk. & M.A. Curtis, Grevillea 3: 51. 1874.

[non. *H. griseum* (Bonord.) Sacc. 1886]

COLONY pinkish. CONIDIOPHORES 159–390 × 3–5 µm. CONIDIOGENOUS CELLS tooth-like. CONIDIA pleurogenous, 15–25-septate, 11.5–15 µm diam., filament width 1–1.5 µm, number of coils 3.5–4 times.

SPECIMENS EXAMINED: BRAZIL. BAHIA: Jeremoabo, on dead fruit, 18.I.2007, A.C.R. Cruz, HUEFS 129350; Senhor do Bonfim, on dead fruit, 28.IX.2006, A.C.R. Cruz, HUEFS 129351; on decaying twig, 28.IX.2006 A.C.R. Cruz, HUEFS 129352. PERNAMBUCO, Buíque, on rotten leaf, 21.VIII.2006 A.C.R. Cruz, HUEFS 129353.

DISTRIBUTION: Australia, Belgium, Brazil, Canada, China, Costa Rica, Cuba, Czech Republic, formerly USSR, Germany, Hungary, Japan, Mexico, New Guinea, New Zealand, Panama, South Africa, Taiwan, USA (CBS 2009, Cooper 2005, Delgado-Rodriguez et al. 2002, Farr & Rossman 2009, Goos 1989, Heredia et al. 1995, Mercado-Sierra et al. 1997, Paulus et al. 2006, Révay 1998, Tsui et al. 2006, Zhao et al. 2007)

Helicosporium griseum is characterized by much branched conidiophores that frequently anastomose. The conidia are smaller in diameter than those cited by Linder (1929), Goos (1989), and Mercado-Sierra (1984) but are similar to

those reported by Zhao et al. (2007). *Helicosporium lumbricopsis* resembles *H. griseum* but is distinguished by the size of the conidia filament, diameter, and stouter conidiophores (Linder 1929, Zhao et al. 2007). This species was collected in Brazil under the name *H. griseum* (Bonord.) Sacc. and deposited at Centraalbureau voor Schimmelcultures (CBS) by R.F. Castañeda Ruiz (CBS 2009). According to Castañeda Ruiz (pers. comm.), the culture produces velvety colonies and anastomosing conidiophores and so is believed to represent *H. griseum* Berk. & M.A. Curtis.

Helicosporium nizamabadense P. Rag. Rao & D. Rao, Mycopathol. Mycol. Appl. 24: 34. 1964.

COLONY white. CONIDIOPHORES 60–155 × 2–3.5 µm. CONIDIOGENOUS CELLS tooth-like. CONIDIA pleurogenous, 15–24 µm diam., filament width 1–3 µm, number of coils 3–4.5 times.

SPECIMENS EXAMINED: BRAZIL. BAHIA: Serra do Ramalho, on decaying leaves of unidentified dicotyledonous plant, 07.I.2008, S.M. Leão-Ferreira, HUEFS 136881.

DISTRIBUTION: Brazil, India (Rao & Rao 1964).

The examined material has a white colony and conidia are produced from distinct denticles on conidiogenous cells (Rao & Rao 1964). *Helicosporium dentophorum* G.Z. Zhao et al., *H. sympodiophorum* G.Z. Zhao et al., and *H. talbotii* Goos (Goos 1989, Zhao et al. 2007) have similar denticles. However, *H. dentophorum* differs by the presence of smaller conidiophores and bigger conidia, *H. sympodiophorum* has bigger conidiophores and conidia, and *H. talbotii* has conidia with narrower filaments. This is the first record for the American continent.

Helicosporium panacheum R.T. Moore, Mycologia 46: 92, 1954.

COLONY white. CONIDIOPHORES 35–135 × 3–4.5 µm. CONIDIOGENOUS CELLS tooth-like. CONIDIA pleurogenous, multiseptate, 12–16 µm diam., filament width 1–1.5 µm, number of coils 2–3.5 times.

SPECIMENS EXAMINED: BRAZIL. BAHIA, Jeremoabo, “Mata das Pororocas”, on dead bark, 16.I.2007, A.C.R. Cruz, HUEFS 129348; Senhor do Bonfim, “fazenda Passaginha”, on dead bark, 5.X.2006, A.C.R. Cruz, HUEFS 129349.

DISTRIBUTION: Brazil, Canada, China, Japan, Mexico, Taiwan, USA, (Arias et al. 2000, CBS 2009, Chang 2001, Goos 1989, Moore 1954, Zhao et al. 2007).

The specimens examined have conidia smaller than in the original description (Moore 1954) but possess the typical characteristics of the species, having hyaline, branched conidiophores at apex and conidia that are white in mass. Zhao et al. (2007) described the same species with wider conidiophores. *Helicosporium panacheum* is reported for the first time in South America.

Helicosporium pannosum (Berk. & M.A. Curtis) R.T. Moore, Mycologia 49: 582. 1957.

TEL.: *Thaxteriella helicoma* (W. Phillips & Plowr.) J.L. Crane, Shearer & M.E. Barr, Can. J. Bot. 76: 610. 1998.

COLONY brown. CONIDIOPHORES $95\text{--}215 \times 6\text{--}7.5 \mu\text{m}$. CONIDIOGENOUS CELLS tooth-like. CONIDIA pleurogenous, 25–50-septate, $34.5\text{--}45 \mu\text{m}$ diam., filament width $3\text{--}4.5 \mu\text{m}$, number of coils 1.5–4 times.

SPECIMENS EXAMINED: BRAZIL. BAHIA, Senhor do Bonfim, “fazenda Passaginha”, on decaying fruit, 5.X.2006, A.C.R. Cruz, HUEFS 129354.

DISTRIBUTION: Argentina, Australia, Brazil, Chile, China, Cuba, Guiana, Haiti Japan, New Zealand, Papua New Guinea, Seychelles Islands, Sri Lanka, Suriname, Tanzania, Thailand, United Kingdom, USA (Barr 1980, CBS 2009, Delgado-Rodríguez et al. 2002, Farr & Rossman 2009, Hughes 1978, Linder 1929, Matsushima 1975, Pinruan et al. 2007, Romero 1983, Sivanesan 1984, Tsui et al. 2001).

Helicosporium pannosum has a very wide widest conidia filament. The conidia of the material examined are smooth, but the specimens examined by Goos (1989) are minutely echinulate. *Helicosporium pannosum* is close to *H. hongkongense* K.M. Tsui et al. with a similar conidia filament width, but its conidiophores are always simple and conidia are subhyaline to yellowish (Tsui et al. 2001). Conidia are smaller than material studied by Zhao et al. (2007).

Helicosporium virescens (Pers.) Sivan., Bitunicate Ascomycetes and their Anamorphs (Vaduz): 591. 1984.

TEL.: *Tubeufia cerea* (Berk. & M.A. Curtis) Höhn., Sber. Akad. Wiss. Wien, Math.-naturw. Kl., Abt. 1, 128: 562. 1919.

COLONY yellow. CONIDIOPHORES $125\text{--}375 \times 3\text{--}6 \mu\text{m}$. CONIDIOGENOUS CELLS tooth-like. CONIDIA pleurogenous, multiseptate, $7.5\text{--}13 \mu\text{m}$ diam., filament width $0.75\text{--}1.2 \mu\text{m}$, number of coils 2–3.5 times.

SPECIMENS EXAMINED: BRAZIL. BAHIA: Campo Formoso, “Mata da Esterzinha”, on decaying twig, 19.IX.2006, A.C.R. Cruz, HUEFS 129355; Jeremoabo, on dead bark, 17.I.2007 A.C.R. Cruz, HUEFS 129356; Senhor do Bonfim, “Carrapichel”, on decaying spathe of *Syagrus coronata* (Mart.) Becc., 9.X.2006, A.C.R. Cruz, HUEFS 129357; Senhor do Bonfim, “Serra da Maravilha”, on dead bark, 11.X.2006, A.C.R. Cruz, HUEFS 129358. PERNAMBUCO, Buíque, on rotten leaf, 11.X.2006, A.C.R. Cruz, HUEFS 129359.

DISTRIBUTION: Austria, Belgium, Brazil, Canada, China, Costa Rica, Cuba, France, Germany, Guiana, Hungary, India, Lithuania, Mexico, Netherlands, Poland, Republic of Belarus, Thailand, United Kingdom, USA (Barr 1980, Farr & Rossman 2009, Goos 1989, Heredia et al. 1995, Linder 1929, Mercado-Sierra et al. 1997, 1998, Révay 1998, Sivanesan 1984, Tsui et al. 2006, Yurchenko 2001, Zhao et al. 2007).

The characteristics observed on the specimens examined are included within the morphological variation of the species (Goos 1989, Zhao et al. 2007). *Helicosporium virescens* differs from *H. aureum* and *H. murinum* Goos by conidiophores that are mostly unbranched. This species was found for the first time for Brazil.

Key to species of *Helicosporium*

1a. Conidia borne acrogenously	2
1b. Conidia borne acropleurogenously	3
1c. Conidia borne pleurogenously	5
2a. (1a). Conidiophores less than 40 µm long, conidia coiled 2–2.25 times	<i>H. dentophorum</i>
2b. Conidiophores more than 40 µm long, conidia coiled 3.5–4 times	<i>H. sympodiophorum</i>
3a. (1b). Conidia more than 20 µm diam.	<i>H. panacheum</i>
3b. Conidia less than 20 µm diam.	4
4a. (3b). Vesicle absent at the apex of conidiophores.	<i>H. gracile</i>
4b. Vesicles usually present at the apex of conidiophores	<i>H. vesiculiferum</i>
5a. (1c). Conidiogenous cells bladder-like.	<i>H. hiospiroides</i>
5b. Conidiogenous cells tooth-like and bladder-like	6
5c. Conidiogenous cells tooth-like	9
6a. (5b). Conidia less than 10 µm diam	<i>H. decumbens</i>
6b. Conidia more than 10 µm diam	7
7a. (6b). Conidiophores mostly unbranched.	<i>H. virescens</i>
7b. Conidiophores mostly branched.	8
8a. (7b). Conidiophores often branched and entangled above.	<i>H. aureum</i>
8b. Conidiophores branched along the mains axis.	<i>H. guianense</i>
9a. (5c). Two kinds of conidiophores (macro and micro)	<i>H. indicum</i>
9b. One kind of conidiophores	10
10a. (9b). Conidiophores consistently branched, anastomosing frequently.	11
10b. Conidiophores simple, sparsely branched, or occasionally anastomosing	12
11a. (10a). Conidiophores subhyaline to dark brown, ascending or more or less erect	<i>H. griseum</i>
11b. Conidiophores dark, erect, at first simple, later branching.	<i>H. lumbricopsis</i>
12a. (10b). Conidiophores mostly colorless, rarely pale brown	13
12b. Conidiophores pale brown to brown	14
13a. (12a). Conidia 10–15 µm diam	<i>H. pallidum</i>
13b. Conidia 18–28 µm diam	<i>H. nizamabadense</i>
14a. (12b). Conidia less than 24 µm diam	15
14b. Conidia mostly more than 24 µm diam.	18
15a. (14a). Conidiophores arising from repent mycelium.	<i>H. talbotii</i>
15b. Conidiophores not as above	16
16a. (15b). Conidial filaments 1–1.5 µm thick	<i>H. murinum</i>
16b. Conidial filaments 1.5–2.7 µm thick	17

- 17a. (16b). Conidiophores mostly simple, rarely branched, conidial filaments
1.5–2 μm thick. *H. abuense*
- 17b. Conidiophores branched from basal parts, conidial filaments
2–2.7 μm thick. *H. phragmitis*
- 18a. (14b). Conidia more than 60 μm diam. *H. raghuveeri*
- 18b. Conidia less than 60 μm diam. 19
- 19a. (18b). Conidiophores simple, conidia subhyaline to yellowish ... *H. hongkongense*
- 19b. Conidiophores simple or branched from basal parts, conidia
subhyaline to pale brown. *H. pannosum*

Acknowledgments

We are deeply indebted to Dr. Guozhu Zhao (Beijing Forestry University, China) and Dr. Eric McKenzie (Landcare Research, New Zealand) for kindly reviewing the manuscript and for many suggestions that greatly improved it. The authors thank Drs. Rafael Castañeda-Ruiz, Gary J. Samuels, Antonio Hernández-Gutiérrez and Carlos Inácio for their generous and valuable assistance with literature. LFP Gusmão and ACR Cruz thank the CNPq (Proc. 474589/2008-0) and “Programa de Pós-graduação em Botânica – PPGBot/UEFS”. ACR Cruz extends thanks to FAPESB (Proc. APR071/2007). Part of the support for this work came from the Program of Research of Biodiversity in the Brazilian Semi-arid (PPBIO Semi-arid/Ministry of Technology and Science).

Literature cited

- Arias RM, Heredia G, Estebanez MR. 2000. Primer registro para México de cinco especies de hongos conidiales helicospóricos. *Rev. Mex. Mic.* 16: 27–32.
- Barr ME. 1980. On the family *Tubeufiaceae* (*Pleosporales*). *Mycotaxon* 12: 137–167.
- Batista AC, Peres GEP, Bezerra JL, Cavalcante W de A. 1964. *Hiospira* Moore e outros hyphomycetes do Amazonas. 364–369, in Anais do XIII Congresso da Sociedade Botânica do Brasil de 1962. Sociedade Botânica do Brasil.
- Berkeley MJ. 1874. Notices of North American fungi. *Grevillea* 3: 49–64.
- Castañeda-Ruiz RF, Gusmão LFP, Heredia-Abarca G, Saikawa M. 2006. Some hyphomycetes from Brazil. Two new species of *Brachydesmiella*, two new combinations for *Repetophragma*, and new records. *Mycotaxon* 95: 261–270.
- Cavalcanti MS, Milanez AI. 2007. Hyphomycetes isolados da água e do solo da Reserva Florestal de Dois Irmãos, Recife, PE, Brasil. *Acta Bot. Bras.* 21: 857–862.
- CBS 2009. Centraalbureau voor schimmelcultures Filamentous fungi database. Centraalbureau voor Schimmelcultures. Utrecht, The Netherlands [<http://www.cbs.knaw.nl/fungi/BioloMICS.aspx>] (viewed online on April 27, 2009)].
- Chang HS. 2001. *Helicoon doliiformis* sp. nov. and two similar helicosporous hyphomycetes from Taiwan. *Bot. Bull. Acad. Sin.* 42: 149–152.
- Cooper JA. 2005. New Zealand hyphomycete fungi: additional records, new species, and notes on interesting collections. *N. Z. J. Bot.* 43: 323–349.
- Cruz ACR, Gusmão LFP, Castañeda-Ruiz RF. 2007a. Conidial fungi from the semi-arid Caatinga biome of Brazil. *Subramaniomyces pulcher* sp. nov. and notes on *Sporidesmium circinophorum*. *Mycotaxon* 102: 25–32.

- Cruz ACR, Gusmão LFP, Leão-Ferreira SM, Castañeda-Ruiz RF. 2007b. Conidial fungi from the semi-arid Caatinga biome of Brazil. *Diplococcium verruculosum* sp. nov. and *Lobatopedis longistriatum* sp. nov. Mycotaxon 102: 33–38.
- Delgado-Rodríguez G, Mena-Portales J, Caldach M, Decock C. 2002. Hyphomycetes (Hongos Mitospóricos) del Área Protegida Mil Cumbres, Cuba Occidental. Crypto. Mycol. 23: 277–293.
- Dix NJ, Webster J. 1995. Fungal Ecology. London, Chapman and Hall.
- Farr DF, Rossman AY. 2009. Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA. [<http://nt.ars-grin.gov/fungaldatabases/>] (viewed online on April 27, 2009)].
- Goos RD. 1980. Some helicosporous fungi from Hawaii. Mycologia 72: 595–610.
- Goos RD. 1985a. On the anamorph genus *Cirrenalia*. Proc. Indian Acad. Sci. (Plant Sci.) 94: 245–252.
- Goos RD. 1985b. A review of the anamorph genus *Helicomycetes*. Mycologia 77: 606–618.
- Goos RD. 1985c. The anamorph genus *Zalerion*. Mycotaxon 23: 445–449.
- Goos RD. 1986. A review of the anamorph genus *Helicoma*. Mycologia 78: 744–761.
- Goos RD. 1987. Fungi with a twist: the helicosporous hyphomycetes. Mycologia 79: 1–22.
- Goos RD. 1989. On the anamorph genera *Helicosporium* and *Drepanospora*. Mycologia 81: 356–374.
- Goos RD. 1990. Review of the anamorph genus *Xenosporium*. Mycologia 82: 742–752.
- Gusmão LFP, Leão-Ferreira SM, Marques MFO, Almeida DAC. 2008. New species and records of *Paliphora* from the Brazilian semi-arid region. Mycologia 100: 306–309.
- Hennings P. 1902. Fungi S. Paulensis II. a cl. Puttemans collecti. Hedwigia 41: 295–311.
- Heredia G, Mercado-Sierra A, Mena-Portales J. 1995. Conidial fungi from leaf litter in a mesophilic cloud forest of Veracruz, Mexico. Mycotaxon 55: 473–490.
- Hughes SJ. 1978. New Zealand fungi 25. Miscellaneous species. N. Z. J. Bot. 16: 311–370.
- Leão-Ferreira SM, Cruz ACR, Castañeda-Ruiz RF, Gusmão LFP. 2008. Conidial fungi from the semi-arid Caatinga biome of Brazil. *Brachysporiellina fecunda* sp. nov. and some new records for Neotropica. Mycotaxon 104: 309–312.
- Linder DH. 1929. A monograph of the Helicosporous Fungi Imperfecti. Ann. Missouri Bot. Gard. 16: 227–388.
- Link HF. 1809. Observations in ordines plantarum naturales. Dissertatio. I. Complectens anadradum ordines epiphytas, Mucedines, Gastromycos et fungos. Ges. Naturf. Freunde Berlin Mag. Neuesten Endec. Gesamtmen Naturk. 3: 3–42.
- Matsushima T. 1975. Icones Microfungorum a Matsushima Lectorum. Kobe, Publ. by author.
- Mendes MAS, Silva VL, Dianese JC, Ferreira JASV, Santos CEN, Gomes-Neto E, Urden AF, Castro C. 1998. Fungos em plantas no Brasil. Brasília, EMBRAPA-SPI.
- Mercado-Sierra A. 1982. Taxonomia y aspectos ecológicos de algunos hifomicetes Helicosporicos Hallados en Cuba. Acta Bot. Cubana 11: 1–11.
- Mercado-Sierra A. 1984. Hifomicetes demaciáceos de Sierra del Rosario, Cuba. Havana, Editorial Academica.
- Mercado-Sierra A, Gené J, Figueras MJ, Rodrigues K, Guarro J. 1998. New or rare hyphomycetes from Cuba. IX. Some species from Pinar del Río Province. Mycotaxon 68: 417–426.
- Mercado-Sierra A, Gené J, Guarro J. 1997. Some Costa Rican Hyphomycetes. II. Mycotaxon 64: 7–15.
- Moore RT. 1953. The North Central *Helicosporae*. Proc. Iowa Acad. Sci. 60: 202–216.
- Moore RT. 1954. Three new species of *Helicosporae*. Mycologia 46: 89–92.
- Moore RT. 1955. Index to the *Helicosporae*. Mycologia 47: 90–103.

- Morgan AP. 1892. North American *Helicosporae*. Jour. Cincinnati Soc. Nat. Hist. 15: 39–52.
- Morris EF. 1978. Belizean hyphomycetes. Mycotaxon 7: 265–274.
- Paulus BC, Kanowski J, Gadek PA, Hyde KD. 2006. Diversity and distribution of saprobic microfungi in leaf litter of an Australian tropical rainforest. Mycol. Res. 110: 1441–1454.
- Pennington RT, Prado DE, Pendry CA. 2000. Neotropical seasonally dry forests and Quaternary vegetation changes. J. Biogeogr. 27: 261–273.
- Pinruan U, Hyde KD, Lumyong S, McKenzie EHC, Jones EBG. 2007. Occurrence of fungi on tissues of the peat swamp palm *Licuala longicalycata*. Fungal Divers. 25: 157–173.
- Prado DE. 2000. Seasonally dry forests of tropical South America: from forgotten ecosystems to a new phytogeographical unity. Edinburgh. J. Bot. 57: 437–461.
- Rao PR, Rao D. 1964. Some *Helicosporae* from Hyderabad. II. Mycopathol. Mycol. Appl. 24: 27–34.
- Révay Á. 1998. Review of the hyphomycetes of Hungary. Stud. Bot. Hung. 27–28: 5–74.
- Romero AI. 1983. Contribución al estudio de los hongos xilófilos de la Argentina. II. Ascomycotina en *Eucalyptus viminalis* (Myrtaceae). Bol. Soc. Argent. Bot. 22: 57–79.
- Romero AI, Pildain MB. 2004. Contribución al estudio de los hongos xilófilos de la Argentina. VII. Ascomycotina en *Eucalyptus viminalis* (Myrtaceae). Bol. Soc. Micol. Madrid 28: 125–134.
- Saccardo PA. 1886. Sylloge Fungorum Omnium Hucusque Cognitorum 4: 1–807.
- Sampaio EVSB. 1995. Overview of the Brazilian Caatinga. 35–58, in SH Bullock, HA Mooney, E Medina (eds.), Seasonally dry forests. Cambridge, Cambridge University Press.
- Samuels GJ, Rossman AY, Müller E. 1978. Life-history studies of Brazilian Ascomycetes. 6. Three species of *Tubeufia* with, respectively, dictyosporous, pycnidial and helicosporous anamorphs. Sydowia 31: 180–192.
- Sivanesan A. 1984. The Bitunicate Ascomycetes and their Anamorphs. Vaduz, Lubrecht & Cramer Ltd.
- Tsui CKM, Berbee ML. 2006. Phylogenetic relationships and convergence of helicosporous fungi inferred from ribosomal DNA sequences. Mol. Phyl. Evol. 39: 587–597.
- Tsui CKM, Goh TK, Hyde KD, Hodgkiss IJ. 2001. New species or records of *Cacumisporium*, *Helicosporium*, *Monotosporella* and *Bahusutrabejia* on submerged wood in Hong Kong streams. Mycologia 93: 389–397.
- Tsui CKM, Sivichai S, Berbee ML. 2006. Molecular systematics of *Helicoma*, *Helicomycetes* and *Helicosporium* and their teleomorphs inferred from rDNA sequences. Mycologia 98: 94–104.
- Yurchenko EO. 2001. On some wood-inhabiting dematiaceous hyphomycetes with remarkable conidia in Belarus. Mycena 1: 32–54.
- Zhao GZ, Liu XZ, Wu WP. 2007. Helicosporous hyphomycetes from China. Fungal Divers. 26: 313–524.