1 2	Oxidative Status of the Ultra-Processed Foods in the
3	Western Diet
4	
5	
6	
7	
8 9	Lisaura Maldonado-Pereira ^{1,2} , Carlo Barnaba ^{3*} , Ilce Gabriela Medina-Meza ^{1,2*}
10	
11	¹ Department of Chemical Engineering and Materials Science, Michigan State
12	University, East Lansing, MI, USA
13 14 15	³ Institute for Quantitative Health Sciences and Engineering, Michigan State University, East Lansing, MI, USA
16	
17	
18	
19	
20	
21	
23	
24	
25	
26	
27	
28	Corresponding Authors
29	Medina-Meza Ilce Gabriela PhD
31	Department of Biosystems and Agricultural Engineering
32	469 Wilson Rd. Room 302C
33	East Lansing, MI
34	<u>ilce@msu.edu</u>
35	Phone: 517-884-1971
36	Paracha Carla DhD
37 38	Dalilava-Callo, FIID Institute for Quantitative Health Sciences and Engineering, Michigan State University
39	Service Road
40	barnabac@msu.edu
41	East Lansing, MI
42	

43 Abstract

44

Ultra-processed foods (UPFs) and their nutritional value have become a trending topic in 45 46 the scientific community because of their increasing demand, and their potentially adverse 47 effects on human health. Besides the poor nutritional value attributed to UPFs, countless studies have also reported the presence of different dietary oxidized substances in these meals. 48 49 DOxS are associated with several chronic diseases such as cardiometabolic diseases (CMD), 50 cancer, diabetes, Parkinson's, and Alzheimer's disease. In this study, a database of DOxS and 51 other dietary metabolites detected in 63 UPFs meals part of the Western diet is reported. 52 Significant differences were found in DOxS and phytosterol contents between ready-to-eat and 53 fast food (FF). Putative biomarkers were suggested for RTE (brassicasterol) and FF (7α-OH and 54 7β -OH), as well as for all 6 different food categories: dairy (brassicasterol), eggs & derivatives 55 (stigmasterol and β -sitosterol), meat & poultry (7 α -OH), seafood, baby food (β -sitosterol), and 56 others (campesterol). Ideally, the use of dietary biomarkers could potentially help in the future to 57 identify in an early stage the presence of different chronic diseases, and even, prevent their 58 development. Nevertheless, an exposure assessment is critical to understand the exposure 59 level of DOxS and their relationship with different chronic diseases.

60

61 Keywords: Dietary Oxysterols, phytosterols, ultra-processed foods, cholesterol oxidation
62 products, lipid oxidation

63 **1. Introduction**

64 Metabolic disorders are worldwide linked to poor dietary habits where Ultra processed foods (UPFs) are emerging as a serious threat to public health.^{1, 2} UPFs are a risk factor for metabolic 65 66 disorders. including but not limited to metabolic and cardiovascular disorders. neurodegeneration, and cancer.³⁻⁵ UPFs have also been correlated with eating addictive-67 68 behaviors because of uncontrollable patterns of seeking and bingeing fat and sugar intake from highly palatable foods, causing long-term changes in the brain.⁶⁻⁸ The adverse effect of long-69 70 term exposure to UPFs is linked to their high caloric content, compared to the lower amount of essential micronutrients, such as minerals and vitamins.⁹ UPFs have become the most 71 72 convenient and accessible food option for developed countries reaching 57% of the American's caloric consumption in 2018,¹⁰ where 52.9% of UPFs are mostly consumed away from home by 73 US youth aged 2-19 years.^{11, 12} 74

75 The USDA's 2020-2025 Dietary Guidelines state that 60% of Americans have one or more preventable chronic diseases related to dietary patterns ¹³. In the last decade, the USDA dietary 76 77 and nutritional guidelines have been highlighting the need for healthy habits; however, these dietary recommendations still need to be met by the population. The most popular UPFs 78 79 consumed in the USA are fast-food (FF) including small, large, and non-chain restaurants, and 80 ready-to-eat (RTE), previously processed products, packed for sale which require minimum preparation at home.¹⁴⁻¹⁶ FF and RTE are highly top preference among children and youth ^{17, 18} 81 82 because of the flavor, attractive design and marketing.^{19, 20}

While is true that UPFs undergo extensive changes during industrial transformation to preserve microbiological safety and stability²¹ including high temperatures, exposure to light, type of cooking, packaging conditions, aging and storage, a caveat is that UPFs embody both an unbalanced ingredient profile as they are mostly constructed from a narrow range of cheap, extracted, refined and fractionated ingredients and detrimental processing that promotes

oxidation of sensible compounds such as lipids and cholesterol,²²⁻²⁶ where changes are difficult
to discriminate. Thus, the need for a reliable biomarker of processing is needed.

90 Dietary oxysterols (DOxS) are oxidized compounds derived from cholesterol oxidation in food, ^{27, 28} which have been proposed as markers of thermal processing of food.^{26, 29, 30} We also 91 92 reported the occurrence of DOxS on infant formulations and baby foods, identifying 7-Ketocholesterol as marker of spray dry in infant formulations.²⁶ DOxS are bioactive lipids known 93 to exert cytotoxic, pro-inflammatory and pro-apoptotic properties.^{27, 31-33} The establishment of 94 95 food processing biomarkers will help to distinguish the extent of processing beyond a gualitative 96 classification. Thus, the systematic mapping of DOxS in the food system is needed to identify levels of food processing, especially in food products within the category of UPFs¹⁰ to better 97 98 assessment of the actual effect of processing on the UPFs nutritional quality.

99 The aim of the study is to fingerprint – for the first time – the content of DOxS and other 100 bioactive lipids on >50 fast food and ready-to-eat meals consumed in the United States. These 101 efforts represent the first step for building a large database that will be widely available to 102 consumers and industry. Importantly, the database allows us to search for markers of food ultra-103 processing and evaluate its impact on the nutritional quality of the Western Diet. We believe that 104 the information presented in this study will enable information regarding unintentional 105 processing-formed compounds which will help to understand the relationship between them and 106 the WD with the purpose of ensuring food nutritional safety and health.

107

108 2. Materials and Methods

109

2.1. Materials, Chemicals and Reagents

Methanol was from Sigma-Aldrich Chloroform was obtained from Omni Solv (Burlington, MA), hexane was purchased from VWR BDH Chemicals (Batavia, IL), 1-butanol and potassium chloride (KCI) from J. T. Baker (Allentown, PA) and diethyl ether was purchased from Fisher

113 Chemical (Pittsburgh, PA). Sodium sulfate anhydrous (Na₂SO₄), Sodium chloride,³⁴ were also 114 purchased from VWR BDH Chemicals. standards of 7 α -hydroxycholesterol (7 α -OH), 7 β -115 hydroxycholesterol (7 β -OH), 5,6 α -epoxycholesterol (5,6 α -epoxy), 5,6 β -epoxycholesterol (5,6 β -116 epoxy), triol, 7-ketocholesterol (7-keto) were purchased from Steraloids (Newport, RI), and 117 purified by using aminopropyl³⁵ cartridges (500 mg/ 3 mL) from Phenomenex (Torrance, CA).

118

2.2. UPFs Selection, Collection and Preparation

The UPFs were selected, and classified into two main groups: fast-foods (FF) and ready-to-119 eat according to our previous study.³⁶ FF (n=23) were collected from retail stores, supermarkets, 120 121 food chains, restaurants, and takeaway in the Greater Lansing area (Michigan, USA) between February 2018 and October 2019, covering 75% of the national market.^{16,17,37} The RTE (n=39) 122 were selected based on the Total Dietary Study 2011-2017 inventory ³⁸ ³⁹. The complete list of 123 the food meals, their respective test code, and their group is provided in Table 1. Additionally, 124 125 foods were grouped into categories according to the fat source as follows: eggs and egg's 126 derivatives (E), dairy products (D), meat and poultry (MP), seafood (S), baby food (BF). Food 127 items that did not fit in any of the previous categories, such as potato-products (potato crisps 128 with and without added flavors, and French fries from restaurants and takeaway, pasta, salad 129 dressings, and popcorn (sweet or salty) were grouped as other products (O).

130 131
 Table 1. Ultra-processed foods meals ID used in this study, divided by group and category.

Category	Sample ID	Ultra-processed Foods (UPF)	Group
	D1 - RTE	American Cheese – Happy Farms	
	D2 - RTE	Cheddar cheese – Happy Farms	
		Margarine (regular, not low-fat, salted) – Countryside	
	DS-RIE	Creamery	
	D5 - RTE	Cream (half & half) -Meijer	Deedute
	D6 - RTE	Swiss cheese - Kroger	Ready to
	D7 - RTE	Cream cheese – Happy Farms	Eat (KIE)
	D8 - RTE	lce cream (regular, not low-fat, vanilla) - Purple Cow	
	D9 - RTE	Yogurt (low-fat, fruit flavored) - Yoplait	
	D10 - RTE	Chocolate milk - Nesquick	
	D11 - RTE	Infant formula – Little Journey (with iron milk-based	

		powder)		
	* D4 – RTE	Butter – Praire Farms		
	MP1 - RTE	Bologna - Eckrich		
	MP2 - RTE	Salami – Oscar Mayer		
		Soup bean w/bacon/pork (canned, prepared w/water) -		
	IVIP3 - KIE	Campbell's		
Meat &	MP4 – RTE	Chili con carne w/beans (canned) - Campbell's		
Poultry	MP5 – RTE	Lasagna w/meat (frozen, heated) – Michael Angelo's		
	MP6 – RTE	Chicken noodle soup - Kroger		
	MP7 – RTE	Beef and vegetables soup – Kroger		
	MP8 – RTE	Mini Ravioli - Chef Boyardee		
	MP9 - RTE	Spaghetti - Chef Boyardee		
Saafaad	C1 DTE	Clam chowder (New England, canned, prep w/ whole		
Sealoou	ST - KIE	milk) – Kroger		
Eggs &	E1 - RTE	Mayonnaise (regular, bottled) – Hellmann's		
derivatives	E2 - RTE	Macaroni salad (from grocery/deli) - Meijer		
	BF1 - RTE	Baby food - beef and broth/gravy – Beech Nut		
	BF2 - RTE	Baby food – chicken and broth/gravy - Gerber		
	BF3 - RTE	Baby food - vegetables and beef - Gerber		
	BF4 - RTE	Baby food - vegetables and chicken - Gerber		
	BF5 - RTE	Baby food - chicken noodle dinner - Gerber		
	BF6 - RTE	Baby food - macaroni, tomato and cheese - Gerber	_	
Baby food	BF7 - RTE	Baby food - turkey and rice - Gerber		
Daby lood	BF8 - RTE	Baby food – turkey and broth/gravy – Beech Nut		
	BF9 - RTE	Baby food – fruit yogurt - Gerber		
	BF10 - RTE	Baby food – chicken with rice - Gerber		
	BF11 - RTE	Baby food – vegetables and turkey - Gerber	_	
	BE12 - RTE	Baby food – macaroni and cheese with vegetables -		
		Gerber	-	
	BF13 - RTE	Pasta pick-ups (cheese ravioli) - Gerber	-	
	O1 - RTE	Popcorn w/butter (microwave) - Kroger	-	
	02 - RTF	Salad dressing (creamy/buttermilk type, regular) – Aldi's		
Others	02 1112	Tuscan Garden		
	O3 - RTE	Macaroni & cheese (boiled) - Kraft		
	04 - RTE	Macaroni & cheese (microwaved) - Kraft		
	MP10 - FF	Hamburger on bun – McDonald's		
	MP11 - FF	Chicken nuggets – McDonald's		
Meat &	MP12 - FF	Cheeseburger on bun – McDonald's	Fast Food	
Poultry	MP13 - FF	Steak tacos w/beans, lettuce, rice and cheese - Chipotle	(FF)	
	MP14 - FF	Cheese and chicken quesadilla - Chipotle		
	MP15 - FF	Chicken burrito w/lettuce, cheese, pico - Chipotle		

	MP16 - FF	Chicken drumstick - KCF
	MP17 - FF	Chicken wing - KFC
	MP18 - FF	Beef w/vegetables - Panda Express
	MP19 - FF	Chicken w/vegetables - Panda Express
	MP20 - FF	Chicken filet - (broiled sandwich) – Chick Fil'A
	MP21 - FF	Roast beef, ham & Provolone - Jimmy Johns
	MP22 - FF	Sliced turkey and bacon - Jimmy Johns
	MP23 - FF	Supreme pizza - Marco's Pizza
	MP24 - FF	Pepperoni pizza, hand tossed - Domino's
Saafaad	S2 - FF	Fish sandwich on bun – McDonald's
Sealoou	S3 - FF	Fried Shrimp – Panda Express
	O5 - FF	French-Fries – McDonald's
	06 - FF	McDonald's Biscuit - Big Breakfast
Othere	07 - FF	McDonald's Hotcakes - Big Breakfast
Others	08 - FF	Biscuit - KFC
	O9 - FF	French Fries - KFC
	010 - FF	Mashed potato - KFC

132 * The only processed (category 3 NOVA) food in this study.

FF were purchased from the selected franchises and brought to the laboratory for immediate analysis. RTE meals were purchased from different local supermarket stores, and immediately brought to the laboratory. Storage conditions were followed according to the label instructions (fresh foods were kept in a fridge at 4°C and frozen meals were kept at -20°C or the temperature indicated in the label).

139 Once the UPF samples arrived at the laboratory, we promptly recorded the following 140 essential information: (1) UPF's name; (2) price; (3) place of purchase, date, and time of 141 collection; (4) type of food (Ready-to-Eat - RTE or Fast-Food - FF); (5) nutritional declaration 142 (including energy, fat, saturated fatty acids, carbohydrates, sugars, fiber, protein, and salt); (6) 143 portion size; (7) list of ingredients; (8) expiration date; and (9) any other pertinent details. For 144 certain UPFs that required additional preparation before analysis, we strictly followed the 145 manufacturer's instructions, which are listed in Table 2. We conducted the analysis using the 146 Sensory Lab kitchen facility at Michigan State University, ensuring accuracy and precision. The

¹³³

- 147 experiments were performed with three biological replicates, and we stored the samples
- appropriately, taking into consideration their specific food matrices.

 Table 2 – Additional cooking preparations according to the label's instructions.

Food Group	Sample	Sample Collection	Cooking Conditions	Lipid Extraction
Seafood	S1-RTE		Microwave: High heat for 2 min and then for 30 s at a time until cooked, stirring each time, about 3 min.	Entire product used
	S2-FF	McDonald's	-	Entire product except for the bun
	MP3-RTE		Microwave: Covered on	
	MP4-RTE		high heat for 2 ½ to 3 min.	-
	MP5-RTE		Oven bake: 25 - 35 min at 400°F.	Entire product
	MP6-RTE		Microwave: Covered, on	used
	MP7-RTE		high heat for 4 to 5 min or until hot.	_
	MP8-RTE		Microwave: 1 minute 30	
	MP9-RTE		seconds or until warm	
Meat &	MP10-FF	McDonald's	-	Entire product except for the bun
Poultry	MP11-FF	McDonald's	-	Entire product used
	MP12-FF	McDonald's	-	Entire product except for the bun
	MP13-FF	Chipotle	-	Entire product except for the tortilla
	MP14-FF	Chipotle	-	Entire product used
	MP15-FF	Chipotle	-	Entire product except for the tortilla

	MP16-FF	KFC	-	Entire product except for the bones
	MP17-FF	KFC	-	Entire product except for bones
	MP20-FF	Chick Fil'A	-	Entire product except for the bun
	MP21-FF	Jimmy John's	-	
	MP22-FF	Jimmy John's	-	
	O1-RTE		Microwave: 2 – 2 ½ min on high heat.	Entire product except for the
	O2-RTE		-	bread
Others	O3-RTE		Boil pan: Stir for 7 to 8 min in boiling water.	
Uners	O4-RTE		Microwave: Uncovered, on high heat 8 to 10 min or until water is absorbed, stirring every 3 min.	Entire product used

- 150 Not Apply
- 151

152 2.3. Lipid Extraction

A cold lipid extraction according to Folch and coworkers⁴⁰ with some modifications 153 154 depending on the food matrix was performed. Thirty grams of sample were homogenized using 155 an Ultra-Turrax® (Tekmar TP 18/10S1 Cincinnati, OH) for 3 min at 300 rpm, Then, placed in a 156 500 mL glass bottle with screwcap with 200 mL of a chloroform:methanol solution (1:1, v/v). 157 Bottle was kept in an oven at 60°C for 20 min before adding 100 mL chloroform. After 2 min of 158 vortex, the content of the bottle was filtered. The filtrate was mixed thoroughly with a 100 mL of 159 1 M KCl solution. Samples were left overnight at 4°C. Then, the lower phase containing lipids 160 was collected and dried at 60°C with a vacuum evaporator at 25 in Hg. Total fat content was determined gravimetrically and fatty acids profiles were previously reported.³⁶ In addition, fat 161 162 content is reported in both units-percentage of total fatty acid weight (% w/w) per 100 g of

163 fresh sample and weight of total fat per serving size (g per serving size)—for nutritional 164 comparison purposes.

165

2.4. Thiobarbituric Acid Reactive Substances (TBARS)

The method modified by Miller ⁴¹, was used to measure lipid oxidation in all UPFs with some modifications. Briefly, 60 mg of fat were weighed into a 10 mL glass test tube with screw cap and the following reagents were added: 100 μ L BHT solution (0.2 mg/mL in water) and 4.9 mL extracting solution (10% TCA in 0.1 M H₂PO₃). Sample blanks were analyzed along with each sample. A standard curve of 0-5 mL TEP solution (10 μ M) was prepared. Test tubes were incubated overnight in the dark at room temperature. TBARS were expressed as μ g malondialdehyde (MDA)/g fat sample.

173

2.5. Total Cholesterol, Tocopherols and Phytosterols Content

A simultaneous isolation method for cholesterol, tocopherols, and phytosterols developed in our laboratory²⁶ was used. 200 mg of lipid were mixed with 50 μ g of 19-hydroxycholesterol and 140 μ g of 5 α -cholestane as internal standards for the determination of phytosterols, tocopherols and total cholesterol, respectively. Subsequently, the sample was mixed with 10 mL of 1 N KOH solution in methanol and left it under gentle agitation and covered from the light for at least 15 h. One-tenth of the unsaponifiable matter was subjected to silvlation.

Sample was mixed with 100 μ L of pyridine and 100 μ L of the Sweeley's reactive mixture (pyridine/hexamethyldisilazane/trimethylchlorosilane, 10:2:1, v/v/v) at 75°C for 45 min, dried under nitrogen stream and dissolved in 1 mL of n-hexane. One microliter of the silylated solution was injected into a gas chromatograph using a fitted with a ZebronTMZB-5HT (Phenomenex, Torrance, CA) capillary column (30 m × 0.25 mm × 0.25 μ m). Oven temperature conditions were set up as follow: 260 to 300°C at 2.5°C/min, then from 300 to 320°C at 8°C/min, and finally hold at 320°C for 1 min. The injector and detector were both set at 320°C. Helium was used as the

187 carrier gas at a flow rate of 54.0 mL/min, the split ratio at 1:50, and pressure constant at 134188 kPa.

189 2.6. DOxS Quantification

The other nine-tenths of the sample was used for the quantification of dietary DOxS. Purification by NH₂-SPE was performed according to Kilvington et al. ²⁶ The enriched fraction was recovered and evaporated to dryness under a nitrogen stream. Subsequently, the purified fraction was silvlated with Sweeley's reactive mixture (at 75 °C for 45 min), dried under a nitrogen stream and dissolved in 1 mL of n-hexane.

195 DOxS acquisition was done using a GCMS-QP2010 SE single quadrupole Shimadzu 196 Corporation (Kyoto, Japan). with one µL of the silvlated sample. DOxS were injected into a 197 single quad GCMS-QP2010 SE with the following conditions: from 250 to 280 °C at 2°C/min, 198 hold at 280 °C for 7 min, and from 280 to 315°C at 1.5 °C/min. Helium was used as a carrier 199 gas (flow rate of 0.37 mL/min); the split ratio was 1:15 and pressure 49.2 KPa. The interface 200 temperature for the GC-MS was 320 °C, with the electron multiplier voltage set at 0 kV and 201 injector at 320 °C. A fused silica column (30 m x 0.25 mm i.d. x 0.25 um thickness) coated with 202 5% phenyl polysyloxane (Phenomenex Zebron ZB-5) and selected ion monitoring (SIM) mode 203 were used.

204 2.7. Statistical Analysis

205 Descriptive statistics were calculated overall, by category, and by food origin. Both mean, 206 and confidence interval (95%) were computed. When comparing RTE vs FF, data did not follow 207 a normal distribution after a density plot analysis and was not homogenous after a residual plot 208 analysis. Therefore, a Matt-Whitney *U*-test was performed, at p < 0.05 significance level. 209 Statistical differences between food categories were evaluated by using the non-parametric 210 Kruskal-Wallis ANOVA by Ranks test, at p < 0.05 significance level. Spearman's correlation 211 across nutritional and non-nutritional variables was also tested. All the statistical analyses were

- computed using Rstudio (Version 1.4.1717 © 2009-2021 RStudio, PBC.). Principal Component
- 213 Analysis was performed in OriginPro (v. 2023, OriginLab) using correlation matrix on normalized
- 214 data.

215

216 3. Results

217 **3.1.** Fat, sterols content, and FAME nutritional indexes in RTE and FF.

We performed quantitative assessment of total fat, cholesterol and phytosterols content in the selected UPFs (**Figure 1**). Fat content in UPFs was statistically similar between FF and RTE when measured on a weight basis; however, if computed according to the serving size, FF showed a significantly higher amount of fat (**Fig. 1A**). As expected, fat-enriched food categories were dairy, and egg and derivatives, whereas seafood categories had the higher fat content when computed on serving basis (**Fig. 1B**).



Figure 1. Total fat (A, B), cholesterol (C), and sterols (D) contents in UPFs by group and category. Box plots represent confidence interval (25-75%) \pm 1SD, line indicates median and white square indicates mean; outliers are also represented as diamonds. When comparing RTE and FF a paired *t*-test was used. The grouping letters in the graph represent distinct categories within the variable under analysis, highlighting significant differences between the groups as determined by the ANOVA, followed by post-hoc Tukey test (p<0.05).

224 Cholesterol and phytosterols are the sterol-like compounds present in UPFs from animal and vegetable origin, respectively. For cholesterol, no significant difference between RTE and 225 FF was observed, and only marginal differences between food categories were observed (Fig. 226 227 1C). FF had a higher sterols content (Fig. 1D); when looking at food categories, meat and egg 228 products showed higher variance in sterols content (Fig. 1D, right panel). As expected, 229 cholesterol was abundant in dairy products and eggs products (category D and E, respectively). Dietary cholesterol from dairy products ranges between 80 mg to 110 g⁴². The lack of vegetable 230 231 oils and other ingredients source of phytosterols, drive the Dairy category to rank 4th in the phytosterols content. Interestingly, Dairy products also were 4th in the DOxS content. Eggs and 232 derivatives category had the highest phytosterols content among all categories and was the 2nd 233 234 highest category in cholesterol content (Fig. 1 C,D). The UPFs comprising the Eggs category, 235 Mayonnaise (E1-RTE) and macaroni salad (E2-RTE) are not solely composed of eggs. They contain a significant amount of soybean oil 43, 44 which is a main source of phytosterols and 236 237 places this category in the first place compared to the rest of the categories. E's position in the 238 cholesterol content assay was expected due to its already known high cholesterol content which ranges from 193-275 mg per egg ⁴⁵⁻⁴⁸. BF have high cholesterol content and a discrete amount 239 240 of phytosterol (Fig. 1C), due to their formulation that includes meat-derivatives as main ingredients, as well as vegetable oils ^{26, 49-51}. Meat and Poultry had lower cholesterol content 241 242 than Dairy and eggs category (Fig. 1C), with a few exceptions. MP6-RTE, MP16-FF, MP17-FF, 243 and BF1-RTE (samples principally made of chicken, and beef) contain the highest cholesterol 244 amounts (Table S2, Supplemental Information). Therefore, the ingredients used during the 245 confection of these meals need to be taken into consideration. The noodles present in the 246 chicken noodle soup (MP6-RTE) are enriched with eggs, the main reason for this meal's high 247 cholesterol content. Interestingly, the ingredient containing cholesterol in the chicken drumstick (MP16-FF) and chicken wing (MP17-FF) samples is whey (10.3 mg cholesterol per cup).⁵² 248

249 Additionally, for MP16-FF and MP17-FF, there's a potential accumulation of these ingredients 250 during the frying step depending on how frequently the frying oil of these fast food is replaced. 251 In our previous study, we characterized the fatty acids composition of both RTE and FF classes ³⁶. Herein, to help further discussion in oxidative status of UPFs, based on our previous data, we 252 computed a series of fatty acids nutritional indexes as reviewed by Chen and Liu.⁵³ (Fig. 2A). 253 254 The PUFA/SFA index is used to determine the impact of fatty acids on cardiovascular health, since a diet rich in PUFA has been linked to lower serum cholesterol levels.⁵³ In the present 255 study, no significant difference in PUFA/SFA index was observed between FF and RTE. The 256 257 indexes of atherogenicity (IA) and thrombogenicity were both proposed in the early '90s and are 258 linked to specific nutritional attributes of fatty acids. These indexes measure the 259 atherogenic/thrombogenic potential of fatty acids, as a ratio between common saturated fatty acids and total unsaturated fatty acids.⁵⁴ We calculate these indices for the UPFs profiled in this 260 261 study (see Material and Methods). RTE had a significantly higher IA and IT index value 262 compared to FF (Fig. 2A); dairy foods were the class that most contributed to the high IA and IT SFA.³⁶ 263 values (Fig. 1B,C), for their load of Finally. the HH 264 (Hypocholesterolemic/Hypercholesterolemic ratio), the HPI (Health-Promoting Index) and the UI 265 (Unsaturated Index) are higher for foods with higher health benefits. In our database, FF scored 266 higher than RTE for all the three indexes (Fig. 2A).

267

3.2. Oxidative status of the UPFs

268 The thiobarbituric acid reactive substances assay (TBARs) is widely used to measure 269 oxidative stress in biological samples⁵⁵ as well as foods.⁵⁶ Lipid peroxidation results in the 270 formation of conjugated dienes, lipid hydroperoxides and degradation products such as alkanes, 271 aldehydes and isoprostanes. The formed hydroperoxides are further decomposed to flavorful 272 secondary oxidation products, which are mainly aldehydes, such as hexanal, 4-hydroxynonenal 273 (HNE), and malondialdehyde (MDA). The TBARs method detects lipid peroxidation products,

274 such as malondialdehyde (MDA), a final product of fatty acids peroxidation. In this study, MDA 275 concentration values were not significantly different between FF and RTE, but significant 276 differences were found between UPFs' categories (Fig. 3A). The highest values for significant 277 amounts of MDA were observed in Meat and Poultry category, followed by Baby Food and 278 Others category (Table S1 Supplemental information). Notability, in several samples, MDA 279 was not detected. It could be possible that second stage oxidation molecules were not either 280 present or the sample was at later stages of the oxidative chain reaction, in which complex and 281 non-reactive species are formed and unable to be detected.

282 We then quantified the total amount of DOxS, which reports cholesterol oxidation. DOxS are 283 cholesteryl-derivatives that accumulate in significant amount within food matrices because of 284 food manufacturing. Importantly, DOxS have known biological effects in both in vitro and animal models.^{57, 58} The FF group showed lower DOxS content (0.15-0.67 µg/serving 95% CI) 285 286 compared to RTE (10.5-20.1 µg/serving 95% CI) (Fig. 3B). As expected, dairy foods contained higher amounts of DOxS because they are rich in cholesterol.^{42, 59, 60} DOxS content was 287 288 positively correlated with price (both absolute and per serving base), as well as fat and sodium 289 content (p < 0.001) (Fig. 3C). Conversely, the negative correlation of those macronutrients with 290 cholesterol content reaffirms that DOxS accumulation is indeed driven by food processing.



Figure 3. Oxidative status of UPFs contents in UPFs by group and category. (A) TBARs values; (B) DOxS amounts; (C) Pearson correlation between oxidative status and selected compositional variables; (D) heat-maps showing clustering of RTE and FF according to oxidative markers and selected classes of lipids; (E) principal components analysis of lipids and sterols, showing clustering of RTE and FF (left), as well as variables loadings (right).When comparing RTE and FF a paired *t*-test was used. The grouping letters in the graph represent distinct categories within the variable under analysis, highlighting significant differences between the groups as determined by the ANOVA, followed by post-hoc Tukey test (p<0.05).

292 Twelve DOxS (7 α -OH, 7 β -OH, 4 β -OH, 5,6 α -epoxy, 5,6 β -epoxy, triol, 6-keto, 7-keto, 20 α -293 OH, 22-OH, 24-OH, and 25-OH) were identified in FF meals and RTE foods. The DOxS content 294 for individual food samples is reported in the Supplemental Material (Table S4-S5 Supplemental material). DOxS content were significantly higher in the O group, followed by 295 296 MP, and S (**Table 1**). 7α -OH was the most abundant DOxS in all UPFs, followed by 7β -OH. 297 Notably, 4β-OH was only detected in FF meals. Also, two side-chain COPs (22-OH and 25-OH) 298 were the most abundant specifically in baby food samples. The heat-map in Fig. 3D highlights 299 compositional differences between FF and RTE, which were better resolved using sparse 300 Partial-Least Square-Discriminant Analysis (sPLS-DA) (Fig. 3E). For the sPLS-DA we included the FAME data from our previous study published somewhere else ³⁶. The multivariate analysis 301 302 showed that RTE are tightly grouped when considering reduced variables featuring lipid and 303 sterols composition. On the other hand, FF were broadly distributed across a dimension 304 comprising several DOxS, but also long-chain FAME and phytosterols. These differences in 305 oxidative load hints that specific compounds may serve as potential biomarkers for processing.

306 3.2.1. Baby Food (BF)

307 Large quantities of phytosterols and cholesterol in BFs meals are intended to fulfill the 308 infant's nutritional needs during this critical period when their body - especially the brain - is 309 under growth and development. For BF, the ingredients used during their confections such as 310 meat and plant-based oils (i.e., sov. coconut, palm, or sunflower oils, etc.) could be beneficial 311 for infants based on their phytosterols and cholesterol specific nutritional needs. However, 312 infants' diet should be closely monitored to avoid uncontrolled accumulation of these 313 compounds that could result in serious health problem such as accelerated atherosclerosis - a 314 consequence of an increased absorption of phytosterols and cholesterol in infants with 315 sitosterolemia ⁶¹, or the development of parenteral nutrition-associated liver disease which is 316 related to a higher intake of phytosterols ⁶².

317 In addition to phytosterols and cholesterol, we found 0.31 mg per serving of DOxS ingested by infants that is obtained through these same ingredients. Potentially, DOxS could negatively 318 319 affect the infant's health in the long run because of their accumulation through this specific diet. 320 Low concentrations of DOxS (1 nM) have been reported in studies of different cardiometabolic, neurodegenerative, inflammatory bowel, ocular diseases, and have been even related to ageing 321 regulation ⁶³⁻⁶⁵. Unfortunately, even though there's abundant information regarding DOxS and 322 323 adverse health effects on adults, little is known about their relationship with infants' health and 324 their diet.

325 Maldonado and coworkers performed a dietary exposure assessment of DOxS in BF and infant formulas (IF) after their consumption throughout the first year of life ⁶⁶ (Maldonado et al, 326 327 2022). They confirmed the dominant presence of cholesterol and phytosterols amounts ingested 328 and absorbed by this specific population group. 5,6 β -epoxycholesterol, 7 β -OH, and 7 α -OH were 329 the highest DOxS exposed through infants' diet with 1,244.4, 191.8, and 165.5 µg/day, 330 respectively; but 5,6β-epoxycholesterol was the most absorbed DOxS with a value of 0.040 mg/kg-day⁶⁶. Further studies focused on DOxS and their effect on infant's health are required to 331 332 address this knowledge gap.

333 3.2.2. Eggs and Its Derivatives

334 For years, the recommended cholesterol daily intake was 300mg/day, and people took close 335 attention to their egg's daily consumption. Recently, this value was eliminated from the 2015-336 2020 Dietary Guidelines for Americans, the 2016 Japanese Dietary Guidelines, and the 2016 Chinese Dietary Guidelines ⁶⁷ because of the lack of evidence of an overall positive association 337 between egg consumption and risk of coronary heart disease ⁶⁸ ^{69, 70}. Even though, dietary 338 339 cholesterol intake raises plasma cholesterol and LDL cholesterol, the effect is not as significant 340 as originally thought, and its consumption also provides other nutrients such as proteins, 341 unsaturated fats, folate, B vitamins, and minerals which are beneficial to human health (Hu, 342 2017).

Eggs and derivatives emerged as the second-highest category with respect to secondary oxidation products (MDA), indicating a higher extent of lipid oxidation in comparison to other categories. Initially, we hypothesized a positive correlation between MDA levels and DOxS (as observed in the Meat and Poultry category, which will be discussed in the next section). However, for the Eggs category, this was not the case, as it exhibited the lowest content of DOxS among all the groups.

349 3.2.3. Meat and poultry (MP)

350 MP had the highest quantities of MDA compared to the other categories. These results were 351 obtained mainly from samples made of pork and chicken. As we mentioned above, MP and E 352 categories showed an opposite association related to DOxS versus MDA results. MP showed a 353 higher DOxS content with high MDA values, while E showed the lowest DOxS content with high 354 MDA values. As we mentioned before, this behavior could be caused by the presence of 355 phytosterols in the sample -MP had significantly lower phytosterols' contents- because of the 356 addition of other ingredients to the whole meal. MDA values are aligned with previous studies ⁷¹⁻ 357 ⁷⁵ that report lipid oxidation changes in food samples after the addition of different ingredients 358 and spices, with considerable amounts of phytochemicals that hinder the lipid oxidation process 359 in the sample.

360 3.2.4. Dairy products

361 Our initial hypothesis suggested that the lower presence of phytosterols in the Eggs 362 category would lead to a higher DOxS content, as these phytosterols may have less antioxidant 363 effect on cholesterol, resulting in the formation of more DOxS. However, our findings 364 contradicted this hypothesis. Interestingly, the processing of dairy products involves subjecting 365 them to very high temperatures, but the duration of these heating steps is considerably shorter 366 (e.g., pasteurization: 17-20 seconds) compared to other cooking processes used for the 367 samples in our study (e.g., frying: 10-15 minutes). Previous research has indicated that when 368 comparing high cooking temperatures with short cooking times against low cooking

temperatures with longer cooking times, there is typically less formation of different DOxS. Hashari and co-workers reported 360.5 ug 7 β -OH/100 g sample of sausage for oven roasting (150°C) for 15min, but 204.3 ug 7 β -OH/100 g sample sausage for microwave cooking (350°C) for 2min.^{76, 77} Moreover, Alina and co-workers detected 2.28 ppm of 5,6 β -Epoxy after oven roasting mutton samples (190°C) for 25 min, but 0.83 ppm of 5,6 β -Epoxy after grilling (230°C) for 20 min.⁷⁸

Lastly, the D category was one of the only two categories where tocopherols could be detected in. It is possible that they can demote certain types of cancer, heart disease, and other chronic ailments.⁷⁹⁻⁸¹ They are commonly found in vegetables and their oils, but traces of tocopherols were found in the D category of this study. Their presence is potentially associated with the use of vegetable colors and other preservatives used during dairy products manufacturing.

381 3.2.5. Seafood

382 Seafood is known as one of the main sources of several essential lipids (ω -3 fatty acids, 383 vitamins-D, and vitamin B12) which are crucial to different biological functions of the human body⁸², and exert nutraceutical properties⁸³. In 2017, the Center of the Plate: Seafood and 384 385 Vegetarian Consumer Trend Report states that 36% of adults are eating more seafood in place of meat.⁸⁴ Even though the seafood category contains the lowest cholesterol and phytosterols 386 amounts, it ranked on the 3rd position of DOxS content (Fig. 2). Results are comparable with 387 previous studies.⁸⁵⁻⁸⁹ The small number of foods used in this category may be the reason for the 388 389 observed results.

390 *3.2.6.* Other products (O)

The "Others" products category ranked 1st in both tocopherols and DOxS contents, as shown in **Fig. 2**. The higher presence of tocopherols in this category can be attributed to the significant use of vegetable oils during the preparation of these meals. Interestingly, while DOxS have been detected in other foods such as biscuits and French fries (especially those from European countries) according to a study in 2002,⁹⁰ no research has reported their content in the same products available in the Western Diet. It is well-established that high cooking temperatures (ranging from 180°C to 200°C) and the specific type of oil used during deep frying of French fries are primary factors that promote the formation of DOxS. These findings highlight the importance of considering both the cooking methods and the choice of ingredients, such as oils, in the context of dietary habits, as they can significantly influence the formation of potentially harmful compounds like DOxS in our food.

402

403

3.4 DOxS as biomarkers of food processing: a preliminary assessment

404 We looked for specific lipid oxidative biomarkers for differentiating food belonging to the FF and 405 RTE categories. To improve discriminatory power, we used a multivariate exploratory ROC 406 analysis to assist the identification of a minimum number of features to use in the model. The 407 model predicts high power discrimination between FF and RTE even when using only to 408 features (AUC = 0.843), whereas the highest power is achieved with 10 features (AUC = 0.949) 409 (Fig. 4A). The accuracy is >75% for all the considered models (Fig. 4B). From the VIP plot (Fig. 410 4C), a few DOxS were selected as top features, including 25-OH, triol and 7-keto, which were 411 previously highlighted in the heatmap (Fig. 3D). Finally, using the first 2 features (25-OH and 412 triol), we achieved sufficient discrimination between FF and RTE, being only 5 food items (three 413 meat products, one baby food and a miscellaneous) assigned to the wrong category.

414 Overall, our results demonstrate clear differences in the oxidative status of FF and RTE,
415 particularly when referring to DOxS.



Figure 4. Biomarker analysis for UPFs. (A) ROC curves based on cross validation performance using a SVM classification method and a built-in SVM feature ranking method. (B) Predictive accuracies with increasing number of features. (C) Top ten significant features with selected frequencies for RTE and FF groups. (D) Predicted class probabilities as a result of average cross-validation for each sample using the best classifier (according to the AUC value). Misclassified samples are labeled.

416

417

418 **4. Discussion**

419 The use of the ambiguous term "ultra-processed foods" in the NOVA classification has given 420 rise to misleading perceptions among consumers and the scientific community regarding the 421 advantages of food preservation technologies. Numerous studies have linked the consumption of UPFs with adverse health outcomes, but there is a lack of quantitative data or biomarkers to 422 423 assess the "level" of processing in these food items. Moreover, the variability in manufacturing 424 processes and the sample matrix further complicate the analysis of food composition and its 425 metabolites. In this study, we present an initial evaluation of the oxidative status of UPFs using 426 an onsite database and quantitative lipidomics focusing on the main lipid oxidative species.

427 **FF and RTE have distinct oxidative signatures.**

428 We previously reported that RTE meals not only have higher fat content compared to FF meals, but also a different FAME composition.³⁶ These differences are due to the extensive formulation 429 430 and ingredients used to produce more palatable and tasty meals in addition to the cooking and 431 preservation methods employed during their manufacturing. In the present study, we confirmed 432 that formulation also affects the sterols composition of UPFs (Fig. 1), and most importantly their 433 oxidative status. Although both groups (FF and RTE) are considered UPFs, there is a clear distinction between these meals due to type of transformation,^{91 enzymatical} source of lipid and 434 435 protein (animal-based vs plant-based), type of preservation technology (e.g., thermal vs 436 nonthermal, wet versus dry), additives addition (e.g., edulcorates, colorants, acidifies). The 437 question that raises is if the differences in composition and manipulation, from both a 438 manufacturing and preparation point of view, will be reflected in major lipid oxidative markers. 439 For instance, although no difference was observed in secondary oxidative products (measured 440 as MDA), FF accumulates more DOxS, which are derived from chemical oxidation of cholesterol 441 (Fig. 2A,B).

442 **DOxS, UPFs, Nutrition and Public Health**

443 The recent alterations to dietary guidelines worldwide stem from emerging research findings. 444 Some of these studies propose that cholesterol alone may not be the sole culprit responsible for 445 chronic diseases like CHD. Instead, they indicate that various other factors, such as overall 446 dietary patterns and their impact on the gut microbiota, play a significant role in the development of human pathological conditions. Recent studies have reported association of specific gut 447 metagenomic species with multiple plasma metabolites.⁹²⁻⁹⁹ Characterization of these 448 449 interactions was stored in an online atlas created by Dekkers and coworkers,¹⁰⁰ expecting to 450 help the scientific community in the process of understanding the effects of the gut microbiota on health It is possible that DOxS could be also influencing changes on the gut microbiota which 451 could end up affecting human health.¹⁰¹ 452

453 *Identification of oxidized biomarkers*

454 Table S6 (Supplemental Information) shows results of the biomarker analysis using MetaboAnalyst package.¹⁰² Brassicasterol was the potential biomarker for RTE meals, while 7a-455 456 OH and 7β-OH were both suggested for FF meals. While both groups fall under the category of 457 UPFs, the distinct production and processing characteristics of each group have a notable 458 influence on the biomarker analysis outcomes. Our findings suggest that brassicasterol and 7α-459 OH + 7 β -OH may potentially serve as biomarkers for ready-to-eat (RTE) and fast-food (FF) 460 meals, respectively. However, we acknowledge the significant impact of the diverse food 461 categories on these results, and further investigations are required to validate these potential 462 biomarkers. **Table S6** also shows potential biomarkers for each individual food category.

463 Maldonado and co-workers⁶⁶ reported 7α -OH as a potential biomarker of the BF category 464 after comparing it against powered and liquid IF (PIF and LIF, respectively) from Kilvington's 465 study.²⁶ In our study, when comparing BF against all other categories, the biomarker analysis 466 pointed to β -sitosterol as a potential biomarker for this specific category. However, we believe 467 that 7α -OH could potentially be a more accurate biomarker, as it reflects the processing 468 differences between these two types of infant foods. On the other hand, the presence of β - 469 sitosterol in these meals may be more closely associated with the use of vegetable oils, which 470 are key ingredients in many of these products. Thus, further investigation is needed to ascertain 471 the most suitable biomarker for distinguishing between these food categories accurately. 472 Further research quantifying DOxS in a broader dataset of UPFs found in the WD is suggested 473 to increase the diet's variability and evaluate any possible changes in the results obtained from 474 the biomarker analysis. This expanded analysis will provide valuable insights into the specific 475 biomarkers that can best differentiate between different categories of UPFs, contributing to a 476 better understanding of their impact on human health.

Consumer awareness of foods and their healthiness is of critical importance,^{103, 104} and 477 478 varies across culture. A pilot study performed in Europe found that Italian and Dutch consumers have weaker negative opinion towards UPF compared to Brazilian consumers.¹⁰³ However, 479 healthiness perception of UPF seems to be linked to heuristic assessment.¹⁰⁴ For instance, the 480 National Health Institute (NIH) recognizes the need for more data to provide insight into 481 personalized foods with the program precision nutrition.¹⁰⁵ A dietary assessment will enable the 482 483 development of an exposure assessment that would become the starting point of a most-484 needed risk assessment to elucidate the association between these food meals and disease risk.¹⁰⁶ Several dietary markers of exposure for many foods have been putatively identified.¹⁰⁷⁻¹¹⁰ 485

486

487 **5.** Conclusion

Our study revealed significant differences in DOxS content between RTE and FF meals. Similarly, the levels of phytosterols exhibited a significant difference between RTE and FF meals. These findings underscore the substantial influence of the parameters and conditions used in the manufacturing processes of these two groups of UPFs. Furthermore, the composition of the food matrix played a pivotal role in determining the presence of these bioactive lipids, as observed when comparing all six food categories. Overall, not all UPFs are deemed to be "unhealthy", a lot of these compounds such as phytosterols and tocopherols are

495 needed to avoid oxidative species that promote development or worsening of human illnesses. 496 However, for RTE meals, there's definitely room for improvement in finding a solution to avoid 497 the excessesive addition of certain nutritional attributes such as sodium, sugar, and fat as part 498 of the current "solutions" for different nutritional and/or safety processes (i.e., microbial 499 reduction, preservation, shel-life extension, etc.). Meanwhile, FF meals seem to be a big source 500 of DOxS. The intake of these toxicological compounds, raises even more health concerns from 501 a chemical and molecular point of view regarding UPFs. Unfortunately, the presence of DOxS in 502 UPFs increse the risk of developmeing of cardiometabolic disorders (CMD), which are one of 503 the most challenging global health issues of the 21st century, recognized by the World Health Organization (WHO),¹¹¹ and the American Society of Endocrinology,¹¹² Oxidative compounds 504 like these cholesterol oxidation products¹¹³, need to be avoided for the sake of the society's 505 506 health.

507 Therefore, we believe that the use of DOxS as biomarkers could potentially help in a future 508 to identify in an early stage the presence of different chronic diseases, and even, prevent their 509 development. Nevertheless, an exposure assessment is critical to understand the exposure 510 level of these toxicological compounds – DOxS – and their relationship with CMD and other 511 chronic diseases.

Until now, no nutritional database of UPFs has been fully developed yet. Maldonado and coworkers already analyzed the nutritional aspects of these same 63 UPFs evaluated in this study.¹¹⁴ Our results aim to expand that database which will serve as our contribution to the scientific community, and general society, in the ongoing quest of elucidating the relationship of DOxS and human health. The creation of a DOxS database of UPFs in WD developed in this study is just the first step to assess and determine the potential health risk caused by the consumption of these UPFs.

519 Acknowledgments:

520 This study was partially funded by the Centre for Research Ingredients Safety (CRIS) of 521 Michigan State University with the GR100229 grant, and the USDA National Institute of Food 522 and Agriculture, Hatch project MICL02526 to I.G.M.M.

- 523524 Conflict of interest
- 525 Authors declare non conflict of interest for this study.

526

527

528 **References**

- 529 (1) Andrade, C. Ultraprocessed Food and Cardiovascular Risk: Estimating the
- 530 Number Needed to Harm in an Unfamiliar Situation. Indian Journal of
- 531 *Psychological Medicine* **2019**, *41* (5), 501-502. DOI:
- 532 10.4103/ijpsym.ijpsym_340_19.
- 533 (2) Lane, M. M.; Davis, J. A.; Beattie, S.; Gomez-Donoso, C.; Loughman, A.;
- 534 O'Neil, A.; Jacka, F.; Berk, M.; Page, R.; Marx, W.; et al. Ultraprocessed food and
- chronic noncommunicable diseases: A systematic review and meta-analysis of
- 43 observational studies. *Obesity Reviews* **2021**, 22 (3). DOI:
- 537 10.1111/obr.13146.
- (3) Abramova, M.; Singh, R. B.; Chibisov, S.; Cornelissen, G.; Takahashi, T.;
- 539 Singh, V.; Pella, D. Chapter 31 Diet and Cancer: A Dysfunction of the Brain. In
- 540 *The Role of Functional Food Security in Global Health*, Academic Press, 2019;
- 541 pp 525-540.
- 542 (4) Jirillo, F.; Jirillo, E.; Magrone, T. Chapter 30 Synbiotics and the Immune
- 543 System. In *Probiotics, Prebiotics, and Synbiotics*, Academic Press, 2016; pp 449-544 458.
- 545 (5) López-Taboada, I.; González-Pardo, H.; Conejo, N. M. Western Diet:
- 546 Implications for Brain Function and Behavior. *Front Psychol* **2020**, *11*, 564413.
- 547 DOI: 10.3389/fpsyg.2020.564413 From NLM.
- (6) Avena, N. M.; Rada, P.; Hoebel, B. G. Sugar and Fat Bingeing Have Notable
- 549 Differences in Addictive-like Behavior. *The Journal of Nutrition* **2009**, *139* (3),
- 550 623-628. DOI: 10.3945/jn.108.097584 (acccessed 11/15/2022).
- 551 (7) Gordon, E. L.; Ariel-Donges, A. H.; Bauman, V.; Merlo, L. J. What Is the
- Evidence for "Food Addiction?" A Systematic Review. *Nutrients* 2018, *10* (4),
 477.
- (8) Lustig, R. H. Ultraprocessed Food: Addictive, Toxic, and Ready for
- 555 Regulation. *Nutrients* **2020**, *12* (11). DOI: 10.3390/nu12113401.
- (9) Monteiro, C. A.; Cannon, G.; Lawrence, M.; Costa Louzada, M. L.; Pereira
- 557 Machado, P. Ultra-processed foods, diet quality, and health using the NOVA
- classification system. Rome, 2019.
- (10) Juul, F.; Martinez-Steele, E.; Parekh, N.; Monteiro, C. A.; Chang, V. W.
- 560 Ultra-processed food consumption and excess weight among US adults. *British*
- *Journal of Nutrition* **2018**, *120* (1), 90-100. DOI: 10.1017/s0007114518001046.
- (11) Urban, L. E.; Weber, J. L.; Heyman, M. B.; Schichtl, R. L.; Verstraete, S.;
- Lowery, N. S.; Das, S. K.; Schleicher, M. M.; Rogers, G.; Economos, C.; et al.
- 564 Energy Contents of Frequently Ordered Restaurant Meals and Comparison with
- 565 Human Energy Requirements and U.S. Department of Agriculture Database
- 566 Information: A Multisite Randomized Study. *Journal of the Academy of Nutrition*
- 567 and Dietetics **2016**, *116* (4), 590-598.e596. DOI: 10.1016/j.jand.2015.11.009
- 568 PubMed.

- (12) Wang, L.; Martínez Steele, E.; Du, M.; Pomeranz, J. L.; O'Connor, L. E.;
- Herrick, K. A.; Luo, H.; Zhang, X.; Mozaffarian, D.; Zhang, F. F. Trends in
- 571 Consumption of Ultraprocessed Foods Among US Youths Aged 2-19 Years,
- 572 1999-2018. *Jama* **2021**, *3*26 (6), 519-530. DOI: 10.1001/jama.2021.10238 From 573 NLM.
- (13) USDA; USDH. *Dietary Guidelines for Americas 2020-2025*. 2020. (accessed
 April 6, 2021).
- 576 (14) Collins. Ready-to-Eat. HarperCollins Publishers: 2022.
- 577 (15) Huang, L.; Hwang, C. A. 16 In-package pasteurization of ready-to-eat meat
- and poultry products. In *Advances in Meat, Poultry and Seafood Packaging*,
 Woodhead Publishing, 2012; pp 437-450.
- (16) Dunford, E. K.; Wu, J.; Wellard-Cole, L.; Watson, W.; Crino, M.; Petersen,
- 581 K.; Neal, B. A comparison of the Health Star Rating system when used for
- restaurant fast foods and packaged foods. *Appetite* **2017**, *117*, 1-8. DOI:
- 583 10.1016/j.appet.2017.06.005.
- 584 (17) Powell, L. M.; Nguyen, B. T. Fast-Food and Full-Service Restaurant
- 585 Consumption Among Children and Adolescents Effect on Energy, Beverage, and 586 Nutrient Intake. *Jama Pediatrics* **2013**, *167*(1), 14-20. DOI:
- 587 10.1001/jamapediatrics.2013.417.
- (18) Dunford, E. K.; Popkin, B. M.; Ng, S. W. Recent Trends in Junk Food Intake in US Children and Adolescents, 2003-2016. *American Journal of Preventive*
- 590 *Medicine* **2020**, *59* (1), 49-58. DOI: 10.1016/j.amepre.2020.01.023.
- 591 (19) Santana, M. O.; Guimarães, J. S.; Leite, F. H. M.; Mais, L. A.; Horta, P. M.;
- Martins, A. P. B.; Claro, R. M. Analysing persuasive marketing of ultra-processed
- foods on Brazilian television. *International Journal of Public Health* **2020**, 65 (7),
- 594 1067-1077.
- (20) Harris, J. L.; Pomeranz, J. L. Infant formula and toddler milk marketing:
- ⁵⁹⁶ opportunities to address harmful practices and improve young children's diets.
- 597 Nutr Rev 2020, 78 (10), 866-883. DOI: 10.1093/nutrit/nuz095 From NLM.
- (21) Cacace, F.; Bottani, E.; Rizzi, A.; Vignali, G. Evaluation of the economic and
- environmental sustainability of high pressure processing of foods. *Innovative*
- 600 Food Science & Emerging Technologies **2020**, 60. DOI:
- 601 10.1016/j.ifset.2019.102281.
- 602 (22) Arneth, W.; Munch, S. Contents of cholesteroloxides in heated meat
- products 1. Possibility of technological influence. *Fleischwirtschaft* 2002, *82* (6),
 109-113.
- 605 (23) Baggio, S. R.; Bragagnolo, N. Cholesterol oxide, cholesterol, total lipid and
- fatty acid contents in processed meat products during storage. *LWT Food*
- 607 Science and Technology **2006**, *39* (5), 513-520. DOI: 10.1016/j.lwt.2005.03.007.
- 608 (24) Dantas, N. M.; Sampaio, G. R.; Ferreira, F. S.; Labre, T. D.; Torres, E.;
- 609 Saldanha, T. Cholesterol Oxidation in Fish and Fish Products. Journal of Food
- 610 Science **2015**, *80* (12), R2627-R2639. DOI: 10.1111/1750-3841.13124.

- 611 (25) Medina-Meza, L. G.; Barnaba, C.; Barbosa-Canovas, G. V. Effects of high
- 612 pressure processing on lipid oxidation: A review. *Innovative Food Science* &
- 613 *Emerging Technologies* **2014**, 22, 1-10. DOI: 10.1016/j.ifset.2013.10.012.
- 614 (26) Kilvington, A.; Barnaba, C.; Rajasekaran, S.; Laurens, M. L. L.; Gabriela
- 615 Medina-Meza, I. Lipid profiling and dietary assessment of infant formulas reveal
- high intakes of major cholesterol oxidative product (7-ketocholesterol). Food
- 617 Chemistry 2021, 129529. DOI: <u>https://doi.org/10.1016/j.foodchem.2021.129529</u>.
- 618 (27) Maldonado-Pereira, L.; Schweiss, M.; Barnaba, C.; Medina-Meza, I. G. The
- role of cholesterol oxidation products in food toxicity. *Food and Chemical Toxicology* 2018, *118*, 908-939. DOI: 10.1016/j.fct.2018.05.059.
- 621 (28) Poli, G.; Iaia, N.; Leoni, V.; Biasi, F. High cholesterol diet, oxysterols and
- their impact on the gut-brain axis. Redox Experimental Medicine 2022, 2022 (1),
- 623 R15-R25.
- 624 (29) Rodriguez-Estrada, M. T.; Garcia-Llatas, G.; Lagarda, M. J. 7-
- 625 Ketocholesterol as marker of cholesterol oxidation in model and food systems:
- when and how. *Biochemical and biophysical research communications* **2014**, 446
- 627 **(3)**, 792-797.
- 628 (30) Leal-Castaneda, E. J.; Inchingolo, R.; Cardenia, V.; Hernandez-Becerra, J.
- A.; Romani, S.; Rodriguez-Estrada, M. T.; Galindo, H. S. G. Effect of Microwave
- 630 Heating on Phytosterol Oxidation. Journal of Agricultural and Food Chemistry
- 631 **2015**, *6*3 (22), 5539-5547. DOI: 10.1021/acs.jafc.5b00961.
- (31) Bjorkhem, I.; Lovgren-Sandblom, A.; Leoni, V.; Meaney, S.; Brodin, L.;
- 633 Salveson, L.; Winge, K.; Palhagen, S.; Svenningsson, P. Oxysterols and
- 634 Parkinson's disease: Evidence that levels of 24S-hydroxycholesterol in
- 635 cerebrospinal fluid correlates with the duration of the disease. *Neuroscience*
- *Letters* **2013**, *555*, 102-105. DOI: 10.1016/j.neulet.2013.09.003.
- (32) Gargiulo, S.; Gamba, P.; Testa, G.; Leonarduzzi, G.; Poli, G. The role of
- 638 oxysterols in vascular ageing. *Journal of Physiology-London* **2016**, *594* (8), 2095-639 2113. DOI: 10.1113/jp271168.
- (33) Lemaire-Ewing, S.; Prunet, C.; Montange, T.; Vejux, A.; Berthier, A.;
- Bessede, G.; Corcos, L.; Gambert, P.; Neel, D.; Lizard, G. Comparison of the
- 642 cytotoxic, pro-oxidant and pro-inflammatory characteristics of different oxysterols.
- 643 *Cell Biol Toxicol* **2005**, *21* (2), 97-114. DOI: 10.1007/s10565-005-0141-2.
- (34) Overholt, M. F.; Mancini, S.; Galloway, H. O.; Preziuso, G.; Dilger, A. C.;
- Boler, D. D. Effects of salt purity on lipid oxidation, sensory characteristics, and
- 646 textural properties of fresh, ground pork patties. *LWT Food Science and*
- 647 *Technology* **2016**, *65*, 890-896. DOI: 10.1016/j.lwt.2015.08.067.
- 648 (35) Liao, P. L.; Cheng, Y. W.; Li, C. H.; Lo, Y. L.; Kang, J. J. Cholesterol-3-beta,
- 5-alpha, 6-beta-triol induced PI3K-Akt-eNOS-dependent cyclooxygenase-2
- expression in endothelial cells. *Toxicology Letters* **2009**, *190* (2), 172-178. DOI:
- 651 10.1016/j.toxlet.2009.07.012.
- (36) Maldonado-Pereira, L.; Barnaba, C.; de Los Campos, G.; Medina-Meza, I.
- 653 G. Evaluation of the nutritional quality of ultra-processed foods (ready to eat +

- 654 fast food): Fatty acids, sugar, and sodium. *J Food Sci* **2022**. DOI: 10.1111/1750-655 3841.16235.
- (37) Tran, A.; Moran, A.; Bleich, S. N. Calorie changes among food items sold in
- US convenience stores and pizza restaurant chains from 2013 to 2017.
- 658 *Preventive medicine reports* **2019**, *15*, 100932.
- (38) Xue, J. P.; Zartarian, V. G.; Liu, S. V.; Geller, A. M. Methyl mercury
- 660 exposure from fish consumption in vulnerable racial/ethnic populations:
- 661 Probabilistic SHEDS-Dietary model analyses using 1999-2006 NHANES and
- 662 1990-2002 TDS data. Science of the Total Environment **2012**, 414, 373-379.
- 663 DOI: 10.1016/j.scitotenv.2011.10.010.
- 664 (39) FDA. Total dietary study.
- 665 <u>http://www.fda.gov/Food/FoodScienceResearch/TotalDietStudy/ucm184232.htm</u>, 666 2014-2017. (accessed 2017 1/20/2017).
- (40) Folch, J.; Lees, M.; Sloane Stanley, G. A simple method for the isolation and
- purification of total lipides from animal tissues. J biol Chem **1957**, 226 (1), 497-
- 669 **509**.
- (41) Miller, D. D. Food chemistry: A laboratory manual; Wiley New York, 1998.
- (42) Lordan, R.; Tsoupras, A.; Mitra, B.; Zabetakis, I. Dairy Fats and
- 672 Cardiovascular Disease: Do We Really Need to be Concerned? *Foods* **2018**, 7
- 673 (3). DOI: 10.3390/foods7030029.
- (43) USDA. Macaroni salad Meijer. USDA, Ed.; 2021.
- (44) USDA. Mayonnaise Hellmann's. USDA, Ed.; 2021.
- (45) Kang, J. W.; Zivkovic, A. M. Are eggs good again? A precision nutrition
- 677 perspective on the effects of eggs on cardiovascular risk, taking into account
- 678 plasma lipid profiles and TMAO. The Journal of Nutritional Biochemistry 2022,
- 679 100, 108906. DOI: <u>https://doi.org/10.1016/j.jnutbio.2021.108906</u>.
- (46) Kralik, G.; Kralik, Z.; Galović, O.; Hanžek, D. Cholesterol Content and Fatty
- Acids Profile in Conventional and Omega-3 Enriched Eggs. *Brazilian Journal of Poultry Science* **2022**, *20*.
- (47) Ceylan, N.; Ciftçi, I.; Mızrak, C.; Kahraman, Z.; Efil, H. Influence of different
- dietary oil sources on performance and fatty acid profile of egg yolk in laying
- hens. Journal of Animal and Feed Sciences **2011**, 20 (1), 71-83, journal article.
- 686 DOI: 10.22358/jafs/66159/2011.
- (48) Spence, J. D.; Jenkins, D. J.; Davignon, J. Dietary cholesterol and egg
- yolks: not for patients at risk of vascular disease. Can J Cardiol 2010, 26 (9),
- e336-339. DOI: 10.1016/s0828-282x(10)70456-6 From NLM.
- 690 (49) Kilvington, A.; Maldonado Pereira, L.; Torres Palacios, C.; Medina Meza,
- I. Phytosterols and their oxidative products in infant formula. *Journal of Food*
- 692 *Process Engineering* **2019**, e13151.
- (50) Küster, A.; Croyal, M.; Moyon, T.; Darmaun, D.; Ouguerram, K.; Ferchaud-
- Roucher, V. Characterization of lipoproteins and associated lipidome in very
- 695 preterm infants: a pilot study. *Pediatric Research* **2022**. DOI: 10.1038/s41390-
- 696 **022-02159-9**.

- 697 (51) Escobar-Sáez, D.; Montero-Jiménez, L.; García-Herrera, P.; Sánchez-Mata,
- 698 M. C. Plant-based drinks for vegetarian or vegan toddlers: Nutritional evaluation
- of commercial products, and review of health benefits and potential concerns.
- 700 *Food Research International* **2022**, *160*, 111646. DOI:
- 701 <u>https://doi.org/10.1016/j.foodres.2022.111646</u>.
- (52) USDA. Beverages, Whey Protein Isolate. USDA, Ed.; 2019.
- (53) Chen, J. P.; Liu, H. B. Nutritional Indices for Assessing Fatty Acids: A Mini-
- Review. International Journal of Molecular Sciences 2020, 21 (16). DOI:
- 705 10.3390/ijms21165695.
- (54) Ulbricht, T.; Southgate, D. Coronary heart disease: seven dietary factors.
 The lancet **1991**, *338* (8773), 985-992.
- (55) Giustarini, D.; Dalle-Donne, I.; Tsikas, D.; Rossi, R. Oxidative stress and
- human diseases: Origin, link, measurement, mechanisms, and biomarkers.
- 710 Critical Reviews in Clinical Laboratory Sciences **2009**, *46* (5-6), 241-281. DOI:
- 711 10.3109/10408360903142326.
- (56) Papastergiadis, A.; Mubiru, E.; Van Langenhove, H.; De Meulenaer, B.
- 713 Malondialdehyde Measurement in Oxidized Foods: Evaluation of the
- 714 Spectrophotometric Thiobarbituric Acid Reactive Substances (TBARS) Test in
- Various Foods. *Journal of Agricultural and Food Chemistry* 2012, *60* (38), 95899594. DOI: 10.1021/jf302451c.
- (57) Griffiths, W. J.; Abdel-Khalik, J.; Hearn, T.; Yutuc, E.; Morgan, A. H.; Wang,
- Y. Q. Current trends in oxysterol research. *Biochemical Society Transactions*
- 719 **2016**, *44*, 652-658. DOI: 10.1042/bst20150255.
- (58) Zmysłowski, A.; Szterk, A. Oxysterols as a biomarker in diseases. *Clin Chim Acta* 2019, *491*, 103-113. DOI: 10.1016/j.cca.2019.01.022.
- (59) Angulo, A. J.; Romera, J. M.; Ramirez, M.; Gil, A. Determination of
- cholesterol oxides in dairy products. Effect of storage conditions. Journal of
- 724 Agricultural and Food Chemistry **1997**, 45 (11), 4318-4323. DOI:
- 725 10.1021/jf970431+.
- (60) Pikul, J.; Rudzińska, M.; Teichert, J.; Lasik, A.; Danków, R.; Przybylski, R.
- 727 Cholesterol oxidation during storage of UHT-treated bovine and caprine milk.
- 728 International Dairy Journal **2013**, 30 (1), 29-32. DOI:
- 729 10.1016/j.idairyj.2012.11.005.
- (61) Yoo, E.-G. Sitosterolemia: a review and update of pathophysiology, clinical
- 731 spectrum, diagnosis, and management. Ann Pediatr Endocrinol Metab 2016, 21
- 732 (1), 7-14. DOI: 10.6065/apem.2016.21.1.7.
- 733 (62) Kurvinen, A.; Nissinen, M. J.; Andersson, S.; Korhonen, P.; Ruuska, T.;
- Taimisto, M.; Kalliomäki, M.; Lehtonen, L.; Sankilampi, U.; Arikoski, P.; et al.
- 735 Parenteral plant sterols and intestinal failure-associated liver disease in
- neonates. *J Pediatr Gastroenterol Nutr* **2012**, *54* (6), 803-811. DOI:
- 737 10.1097/MPG.0b013e3182474118 From NLM.

- (63) Poli, G.; Biasi, F.; Leonarduzzi, G. Oxysterols in the pathogenesis of major
- chronic diseases. *Redox Biology* **2013**, *1* (1), 125-130. DOI:
- 740 10.1016/j.redox.2012.12.001.
- (64) de Medina, P.; Silvente-Poirot, S.; Poirot, M. Oxysterols are potential
- physiological regulators of ageing. Ageing Res Rev 2022, 77, 101615. DOI:
- 743 10.1016/j.arr.2022.101615.
- (65) Kulig, W.; Cwiklik, L.; Jurkiewicz, P.; Rog, T.; Vattulainen, I. Cholesterol
- oxidation products and their biological importance. *Chem Phys Lipids* 2016, *199*,
 144-160. DOI: 10.1016/j.chemphyslip.2016.03.001.
- (66) Maldonado-Pereira, L.; Barnaba, C.; Medina-Meza, I. G. Dietary exposure
- assessment of infant formula and baby foods? oxidized lipids in the US
- population. *Food and Chemical Toxicology* **2023**, *17*2. DOI:
- 750 10.1016/j.fct.2022.113552.
- (67) Zhu, H.; Chen, Z.-Y. Do We No Longer Need To Worry about Dietary
- 752 Cholesterol? Journal of Agricultural and Food Chemistry 2017, 65 (46), 9931-
- 753 9933. DOI: 10.1021/acs.jafc.7b04829.
- (68) Orsavova, J.; Misurcova, L.; Ambrozova, J.; Vicha, R.; Mlcek, J. Fatty Acids
- 755 Composition of Vegetable Oils and Its Contribution to Dietary Energy Intake and
- 756 Dependence of Cardiovascular Mortality on Dietary Intake of Fatty Acids.
- *International Journal of Molecular Sciences* 2015, *16* (6), 12871-12890. DOI:
 10.3390/ijms160612871.
- (69) Hu, F. B. Diet and Heart Disease. In International Encyclopedia of Public
- 760 *Health*, 2nd ed.; Inc., E. Ed.; 2017; Vol. 2.
- 761 (70) Alphonse, P. A.; Ramprasath, V.; Jones, P. J. Effect of dietary cholesterol
- and plant sterol consumption on plasma lipid responsiveness and cholesterol
- trafficking in healthy individuals. Br J Nutr 2017, 117 (1), 56-66. DOI:
- 764 10.1017/S0007114516004530.
- 765 (71) Jung, S.; Nam, K. C.; Jo, C. Detection of malondialdehyde in processed
- meat products without interference from the ingredients. *Food Chem* **2016**, *209*,
- 767 90-94. DOI: 10.1016/j.foodchem.2016.04.035.
- 768 (72) Wang, W.; Zhang, Z.; Liu, X.; Cao, X.; Wang, L.; Ding, Y.; Zhou, X. An
- 769 Improved GC-MS Method for Malondialdehyde (MDA) Detection: Avoiding the
- ⁷⁷⁰ Effects of Nitrite in Foods. *Foods* **2022**, *11* (9). DOI: 10.3390/foods11091176.
- 771 (73) Sobral, M. M. C.; Casal, S.; Faria, M. A.; Cunha, S. C.; Ferreira, I. M. P. L.
- Influence of culinary practices on protein and lipid oxidation of chicken meat
- burgers during cooking and in vitro gastrointestinal digestion. *Food Chem Toxicol*
- 774 **2020**, *141*, 111401. DOI: 10.1016/j.fct.2020.111401.
- 775 (74) dos Santos, J. J. A.; Pascoal, L. A. F.; Grisi, C. V. B.; Santos, V. d. C.; de
- Santana Neto, D. C.; Jordão Filho, J.; Herminio, M. P. F.; Dantas, A. F. Soybean
- oil and selenium yeast levels in the diet of rabbits on performance, fatty acid
- profile, enzyme activity and oxidative stability of meat. *Livestock Science* **2022**,
- 779 263, 105021. DOI: <u>https://doi.org/10.1016/j.livsci.2022.105021</u>.

- 780 (75) He, W.-S.; Sun, Y.; Li, Z.; Yang, H.; Li, J.; Wang, Q.; Tan, C.; Zou, B.
- 781 Enhanced antioxidant capacity of lipoic acid in different food systems through
- lipase-mediated esterification with phytosterols. Journal of the Science of Food
- 783 and Agriculture **2022**, *n/a* (n/a), <u>https://doi.org/10.1002/jsfa.12073</u>. DOI:
- 784 <u>https://doi.org/10.1002/jsfa.12073</u> (acccessed 2022/10/09).
- (76) Hashari, S. Z.; Rahim, A. A.; Meng, G. Y.; Ramiah, S. K. Quantification of
- Cooking Method Effect on COP Content in Meat Types Using Triple Quadrupole
 GC-MS/MS. *Molecules* 2020, *25* (21). DOI: 10.3390/molecules25214978.
- 788 (77) Khan, M. I.; Min, J. S.; Lee, S. O.; Yim, D. G.; Seol, K. H.; Lee, M.; Jo, C.
- 789 Cooking, storage, and reheating effect on the formation of cholesterol oxidation
- products in processed meat products. *Lipids Health Dis* **2015**, *14*, 89. DOI:
- 791 10.1186/s12944-015-0091-5.
- (78) Rahim, A.; Nurul, A.; Sakinah, F.; Zam Hashari, S.; Juhana, M.; Mashitoh,
- A.; Syuhada, H.; Mawaddah, A.; Nurulhuda, S. Effect of Grilling and Roasting on
- Formation of Cholesterol Oxidation Products (COPs) in Chicken and Mutton.
- 795 World Applied Sciences Journal **2012**, 17, 21-24.
- (79) Shahidi, F.; de Camargo, A. C. Tocopherols and Tocotrienols in Common
- and Emerging Dietary Sources: Occurrence, Applications, and Health Benefits.
 Int J Mol Sci 2016, *17* (10). DOI: 10.3390/ijms17101745.
- (80) Wen Lee, H.; Bi, X.; Jeyakumar Henry, C. Carotenoids, tocopherols and
- 800 phylloquinone content of 26 green leafy vegetables commonly consumed in
- 801 Southeast Asia. *Food Chem* **2022**, *385*, 132729. DOI:
- 802 10.1016/j.foodchem.2022.132729.
- 803 (81) Halder, M.; Petsophonsakul, P.; Akbulut, A. C.; Pavlic, A.; Bohan, F.;
- Anderson, E.; Maresz, K.; Kramann, R.; Schurgers, L. Vitamin K: Double Bonds
- 805 beyond Coagulation Insights into Differences between Vitamin K1 and K2 in
- 806 Health and Disease. Int J Mol Sci 2019, 20 (4). DOI: 10.3390/ijms20040896.
- 807 (82) Hosomi, R.; Yoshida, M.; Fukunaga, K. Seafood consumption and
- components for health. *Glob J Health Sci* **2012**, *4* (3), 72-86. DOI:
- 809 10.5539/gjhs.v4n3p72 From NLM.
- 810 (83) Nag, M.; Lahiri, D.; Dey, A.; Sarkar, T.; Pati, S.; Joshi, S.; Bunawan, H.;
- Mohammed, A.; Edinur, H. A.; Ghosh, S.; et al. Seafood Discards: A Potent
- 812 Source of Enzymes and Biomacromolecules With Nutritional and Nutraceutical
- 813 Significance. *Front Nutr* **2022**, *9*, 879929. DOI: 10.3389/fnut.2022.879929.
- 814 (84) Consumer & Restaurant Menu Trends: The Clean Label Influence;
- 815 Technomic, I., 2020. Retrieved from Healthy eating consumer trend report,
- 816 Technomic. <u>https://cleanlabel.globalfoodforums.com/wp-</u>
- 817 <u>content/uploads/sites/3/2020/08/Consumer-Restaurant-Menu-Trends-Clean-</u>
- 818 <u>Label-L-Freier-A-Harvey-Technomic-2017.pdf</u>.
- (85) Osada; Kyoichi; Takehiro, K.; Li, C.; Koji, Y.; Michihiro, S. Levels and
- formation of oxidized cholesterol in processed marine foods. **1993**, *41*.

- 821 (86) Ohshima, T.; Li, N.; Koizumi, C. Oxidative decomposition of cholesterol in
- fish products. Journal of the American Oil Chemists' Society **1993**, 70 (6), 595-
- 823 600, journal article. DOI: 10.1007/bf02545326.
- 824 (87) Shozen, K.-i. Formation of Cholesterol Oxides in Marine Fish Products
- 825 Induced by Grilling. *Fisheries science* **1995**, *61* (5), 817-821.
- 826 (88) Dantas, N. M.; Sampaio, G. R.; Ferreira, F. S.; Labre, T. a. S.; Torres, E. A.;
- 827 Saldanha, T. Cholesterol Oxidation in Fish and Fish Products. *J Food Sci* **2015**,
- 828 80 (12), R2627-2639. DOI: 10.1111/1750-3841.13124.
- 829 (89) Dantas, N. M.; de Oliveira, V. S.; Sampaio, G. R.; Chrysostomo, Y. S. K.;
- 830 Chávez, D. W. H.; Gamallo, O. D.; Sawaya, A. C. H. F.; Torres, E. A. F. D.;
- 831 Saldanha, T. Lipid profile and high contents of cholesterol oxidation products
- (COPs) in different commercial brands of canned tuna. *Food Chem* **2021**, 352,
- 833 129334. DOI: 10.1016/j.foodchem.2021.129334.
- 834 (90) Guardiola, F. Cholesterol and phytosterol oxidation products: analysis,
- 835 occurrence, and biological effects; The American Oil Chemists Society, 2002.
- 836 (91) McClements, D. J.; Decker, E. A.; Weiss, J. Emulsion-based delivery
- systems for lipophilic bioactive components. *J Food Sci* **2007**, 72 (8), R109-124.
- 838 DOI: 10.1111/j.1750-3841.2007.00507.x From NIm.
- (92) Pedersen, H. K.; Gudmundsdottir, V.; Nielsen, H. B.; Hyotylainen, T.;
- Nielsen, T.; Jensen, B. A.; Forslund, K.; Hildebrand, F.; Prifti, E.; Falony, G.; et
- al. Human gut microbes impact host serum metabolome and insulin sensitivity.
- 842 *Nature* **2016**, *535* (7612), *376-381*. DOI: 10.1038/nature18646.
- 843 (93) Kurilshikov, A.; van den Munckhof, I. C. L.; Chen, L.; Bonder, M. J.; Schraa,
- K.; Rutten, J. H. W.; Riksen, N. P.; de Graaf, J.; Oosting, M.; Sanna, S.; et al. Gut
- 845 Microbial Associations to Plasma Metabolites Linked to Cardiovascular
- 846 Phenotypes and Risk. Circ Res 2019, 124 (12), 1808-1820. DOI:
- 847 10.1161/CIRCRESAHA.118.314642.
- 848 (94) Visconti, A.; Le Roy, C. I.; Rosa, F.; Rossi, N.; Martin, T. C.; Mohney, R. P.;
- Li, W.; de Rinaldis, E.; Bell, J. T.; Venter, J. C.; et al. Interplay between the
- human gut microbiome and host metabolism. *Nat Commun* **2019**, *10* (1), 4505.
- 851 DOI: 10.1038/s41467-019-12476-z.
- (95) Vojinovic, D.; Radjabzadeh, D.; Kurilshikov, A.; Amin, N.; Wijmenga, C.;
- 853 Franke, L.; Ikram, M. A.; Uitterlinden, A. G.; Zhernakova, A.; Fu, J.; et al.
- 854 Relationship between gut microbiota and circulating metabolites in population-
- based cohorts. *Nat Commun* **2019**, *10* (1), 5813. DOI: 10.1038/s41467-019-
- 856 13721-1.
- (96) Bar, N.; Korem, T.; Weissbrod, O.; Zeevi, D.; Rothschild, D.; Leviatan, S.;
- Kosower, N.; Lotan-Pompan, M.; Weinberger, A.; Le Roy, C. I.; et al. A reference
- map of potential determinants for the human serum metabolome. *Nature* **2020**,
- 860 588 (7836), 135-140. DOI: 10.1038/s41586-020-2896-2.
- 861 (97) Partula, V.; Deschasaux-Tanguy, M.; Mondot, S.; Victor-Bala, A.;
- Bouchemal, N.; Lécuyer, L.; Bobin-Dubigeon, C.; Torres, M. J.; Kesse-Guyot, E.;
- 863 Charbit, B.; et al. Associations between untargeted plasma metabolomic

- signatures and gut microbiota composition in the Milieu Intérieur population of
- healthy adults. British Journal of Nutrition **2021**, 126 (7), 982-992. DOI:
- 10.1017/S0007114520004870 From Cambridge University Press Cambridge
 Core.
- (98) Asnicar, F.; Berry, S. E.; Valdes, A. M.; Nguyen, L. H.; Piccinno, G.; Drew,
- D. A.; Leeming, E.; Gibson, R.; Le Roy, C.; Khatib, H. A.; et al. Microbiome
- 870 connections with host metabolism and habitual diet from 1,098 deeply
- phenotyped individuals. *Nat Med* 2021, 27 (2), 321-332. DOI: 10.1038/s41591020-01183-8.
- 873 (99) Forslund, S. K.; Chakaroun, R.; Zimmermann-Kogadeeva, M.; Markó, L.;
- Aron-Wisnewsky, J.; Nielsen, T.; Moitinho-Silva, L.; Schmidt, T. S. B.; Falony, G.;
- Vieira-Silva, S.; et al. Combinatorial, additive and dose-dependent drug-
- microbiome associations. *Nature* **2021**, *600* (7889), 500-505. DOI:
- 877 10.1038/s41586-021-04177-9.
- (100) Dekkers, K. F.; Sayols-Baixeras, S.; Baldanzi, G.; Nowak, C.; Hammar, U.;
- Nguyen, D.; Varotsis, G.; Brunkwall, L.; Nielsen, N.; Eklund, A. C.; et al. An
- online atlas of human plasma metabolite signatures of gut microbiome
- composition. *Nat Commun* **2022**, *13* (1), 5370. DOI: 10.1038/s41467-022-330500.
- (101) Yan, C.; Kwek, E.; Ding, H. F.; He, Z. Y.; Ma, K. Y.; Zhu, H. Y.; Chen, Z. Y.
- 884 Dietary Oxidized Cholesterol Aggravates Chemically Induced Murine Colon
- Inflammation and Alters Gut Microbial Ecology. *Journal of Agricultural and Food Chemistry* 2022. DOI: 10.1021/acs.jafc.2c05001.
- (102) Pang, Z. Q.; Zhou, G. Y.; Ewald, J.; Chang, L.; Hacariz, O.; Basu, N.; Xia,
- J. G. Using MetaboAnalyst 5.0 for LC-HRMS spectra processing, multi-omics
- integration and covariate adjustment of global metabolomics data. *Nature*
- 890 Protocols **2022**, *17* (8), 1735-1761. DOI: 10.1038/s41596-022-00710-w.
- (103) Bolhuis, D.; Mosca, A. C.; Pellegrini, N. Consumer Awareness of the
- 892 Degree of Industrial Food Processing and the Association with Healthiness-A
- 893 Pilot Study. *Nutrients* **2022**, *14* (20). DOI: 10.3390/nu14204438.
- (104) Machin, L.; Antunez, L.; Curutchet, M. R.; Ares, G. The heuristics that
- guide healthiness perception of ultra-processed foods: a qualitative exploration.
- 896 Public Health Nutrition **2020**, 23 (16), 2932-2940. DOI:
- 897 10.1017/s1368980020003158.
- (105) NIH. Nutrition for Precision Health, powered by the All of Us Research
- 899 *Program.* 2022. <u>https://commonfund.nih.gov/nutritionforprecisionhealth</u>
- 900 (accessed.
- 901 (106) Yin, X. F.; Gibbons, H.; Rundle, M.; Frost, G.; McNulty, B. A.; Nugent, A.
- 902 P.; Walton, J.; Flynn, A.; Gibney, M. J.; Brennan, L. Estimation of Chicken Intake
- by Adults Using Metabolomics-Derived Markers. *Journal of Nutrition* **2017**, *147*
- 904 (10), 1850-1857. DOI: 10.3945/jn.117.252197.

- 905 (107) Cross, A. J.; Major, J. M.; Sinha, R. Urinary Biomarkers of Meat
- 906 Consumption. Cancer Epidemiology Biomarkers & Prevention 2011, 20 (6),
- 907 1107-1111. DOI: 10.1158/1055-9965.epi-11-0048.
- 908 (108) O'Gorman, A.; Gibbons, H.; Brennan, L. Metabolomics in the identification
- of biomarkers of dietary intake. *Computational and Structural Biotechnology Journal* 2013, 4 (5). DOI: 10.5936/csbj.201301004.
- 911 (109) O'Gorman, A.; Brennan, L. The role of metabolomics in determination of
- new dietary biomarkers. *Proceedings of the Nutrition Society* **2017**, *76* (3), 295-
- 913 302. DOI: 10.1017/s0029665116002974.
- 914 (110) Prendiville, O.; Walton, J.; Flynn, A.; Nugent, A. P.; McNulty, B. A.;
- 915 Brennan, L. Classifying Individuals Into a Dietary Pattern Based on Metabolomic
- 916 Data. *Molecular Nutrition & Food Research* **2021**, 65 (11). DOI:
- 917 10.1002/mnfr.202001183.
- 918 (111) Organization, W. H. *Cardiovascular diseases*.
- 919 http://www.who.int/mediacentre/factsheets/fs317/en/, 2017. (accessed.
- 920 (112) Rakhra, V.; Galappaththy, S. L.; Bulchandani, S.; Cabandugama, P. K.
- 921 Obesity and the Western Diet: How We Got Here. Mo Med 2020, 117 (6), 536-
- 922 538. From NLM.
- 923 (113) Maldonado-Pereira, L.; Barnaba, C.; Medina-Meza, I. G. Dietary Exposure
- to Oxidized Lipids in US Infants. *medRxiv* **2022**, 2022.2006.2022.22276754. DOI:
- 925 10.1101/2022.06.22.22276754.
- 926 (114) Maldonado-Pereira, L.; Barnaba, C.; De los Campos, G.; Medina-Meza, I.
- 927 Evaluation of the Nutritional Quality of Ultra-Processed Foods (Ready to Eat +
- 928 Fast Food): Fatty Acid Composition. *medRxiv* **2021**, 2021.2004.2016.21255610.
- 929 DOI: 10.1101/2021.04.16.21255610.
- 930
- 931