



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 2, 2015

Mr. Michael Kellogg
Agent for AgraCity Crops and Nutrition, Inc.
c/o Pyxis Regulatory Consulting, Inc.
4110 136th St. NW
Gig Harbor, WA 98332

Subject: Label Amendment – Revised label
Product Name: MPower Propiconazole
EPA Registration Number: 91097-9
Application Date: 09/03/2015
Decision Number: 505279

Dear Mr. Kellogg:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 91097-9
Decision No. 505279

with FIFRA section 6. If you have any questions, please contact Lisa Pahel by phone at 703-347-0459, or via email at pahel.lisa@epa.gov.



Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Enclosure

ACCEPTED

Nov 02, 2015

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 91097-9

MPower Propiconazole

GROUP **3** FUNGICIDE

Broad spectrum fungicide for control of plant, turf and ornamental diseases

ACTIVE INGREDIENT:

Propiconazole* 41.8%

OTHER INGREDIENTS:** 58.2%

TOTAL: 100.0%

*CAS No. 60207-90-1

**Contains petroleum distillate.

Contains 3.6 lbs. propiconazole a.i. per gallon

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Do not give any liquid to the person.• Do not induce vomiting unless told to by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.	

EPA Reg. No. 91097-9

EPA Est. No. XXXXX-XX-XXX

Manufactured For:

AgraCity Crops & Nutrition Inc.
375 E. Horsetooth Rd.
Building 5, Suite 202
Fort Collins, CO 80525

Net Contents:

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

Causes substantial but temporary eye injury. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid breathing spray mist. This product may cause sensitization reactions in certain individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are polyethylene or polyvinyl chloride.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to inter-tidal areas below mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any

requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate or Viton
- Shoes plus socks
- Protective eyewear

PRODUCT INFORMATION

MPower Propiconazole is a broad spectrum fungicide for the control of specified diseases in labeled crops.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

USE RESTRICTION

Do not use this product in greenhouses or as a tree injection.

Rotational Crops

Alfalfa can be planted 75 days after the last MPower Propiconazole application if the total application of propiconazole has not exceeded 0.22 lb. active ingredient per acre during the previous year. Do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of MPower Propiconazole application to the preceding crop unless the second crop appears on this label.

Integrated Pest Management

MPower Propiconazole should be integrated into an overall disease and pest management strategy (IPM) whenever the use of a fungicide is required. Cultural

practices known to reduce disease development should be followed. Consult your local agricultural authorities for additional IPM strategies established for your area. MPower Propiconazole may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

Group	3	Fungicide
--------------	----------	------------------

MPower Propiconazole is in the Group 3 class fungicides. The mode of action of MPower Propiconazole is as a demethylation inhibitor of sterol biosynthesis (DMI) which disrupts membrane synthesis by blocking demethylation. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank mixing with products having different modes of action or limiting the total number of applications per season. AgraCity Crops & Nutrition Inc. encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label. MPower Propiconazole should not be alternated or tank mixed with any fungicide to which resistance has already developed.

SPRAY EQUIPMENT

Thorough coverage is necessary to provide good disease control.

To avoid spray drift, do not apply when conditions favor drift beyond the target areas. Avoid spray overlap, as crop injury may occur.

Air assisted or air blast sprayers move spray droplets into the canopy using a forced air stream. Set up the fan to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate sprayer before use.

Use a pump with capacity to: (1) maintain 35-40 psi at nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension – this requires recirculation of 10% of tank volume per minute. Use a jet agitator or liquid sparge tube for agitation. Do not air sparge.

Although MPower Propiconazole is an emulsifiable concentrate, it is suggested that screens be used to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturer's and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

MIXING INSTRUCTIONS

Prepare no more spray mixture than is required for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

MPower Propiconazole Alone: Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank. With the agitator running, add the MPower Propiconazole to the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the MPower Propiconazole has completely dispersed into the mix water. Maintain agitation until all of the mixture has been sprayed.

MPower Propiconazole + Tank Mixtures: MPower Propiconazole is usually compatible with all tank mix partners listed on this label. To determine the physical compatibility of MPower Propiconazole with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water dispersible granular products first, the liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank. With the agitator running, add the tank mix partner into the tank. Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and the MPower Propiconazole to the spray tank. Allow the MPower Propiconazole to completely disperse. Spray the mixture with the agitator running.

If using MPower Propiconazole in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate may be exceeded, and the most restrictive label precautions and limitations must be followed. This product may not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Do not mix MPower Propiconazole with Syllit[®] or crop injury may result.

APPLICATIONS INSTRUCTIONS

Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply in a manner that will result in exposure to humans or animals.

Ground Application: For tree crops, apply in a minimum of 50 gallons of water per acre unless specified otherwise. For other crops, apply a minimum of 10 gallons of water per acre unless specified otherwise.

Aerial Application: For tree crops, apply in a minimum of 10 gallons of water per acre unless specified otherwise. For other crops, apply in a minimum of 2 gallons of water per acre unless specified otherwise.

Application Through Irrigation Systems (Chemigation): Apply MPower Propiconazole through irrigation equipment only to crops for which chemigation is specified on this label.

For chemigation, apply 0.1-0.25 inches of water. Chemigation with excessive water may lead to a decrease in efficacy.

MPower Propiconazole, alone or in combination with other pesticides which are registered for application through irrigation systems, may be applied through irrigation systems. Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. Do not connect an irrigation system (including green house systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Restrictions: Do not inject MPower Propiconazole at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part MPower Propiconazole. MPower Propiconazole is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used, but should be replaced once a year. Do not use Viton, Buna-N, Neoprene, or PVC seals.

Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended.

Center Pivot Irrigation Equipment

Restrictions: (1) Use only with drive systems which provide uniform water distribution.
(2) Do not use end guns when chemigating MPower Propiconazole through center pivot systems because of non-uniform application.

- Determine the size of the area to be treated.
- Determine the time required to apply 1/8-1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. When applying MPower Propiconazole through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of MPower Propiconazole required to treat the area covered by the irrigation system.
- Add the required amount of MPower Propiconazole and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the MPower Propiconazole solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the MPower Propiconazole solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying MPower Propiconazole through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of MPower Propiconazole required to treat the area covered by the irrigation system.
- Add the required amount of MPower Propiconazole into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop the injection equipment after treatment is completed. Continue to operate the system until the MPower Propiconazole solution has cleared the last sprinkler head.

APPLICATION INSTRUCTIONS

Crop	Pests Controlled	Application Rate/Acre	Instructions
Almonds	Brown Rot Blossom Blight (<i>Monilinia laxa</i> , <i>M. fructicola</i>)	4 - 8 fl. oz.	Apply MPower Propiconazole in at least 15 gallons of spray per acre at 5 - 10% bloom and 50 - 100% bloom using ground or air equipment in sufficient volume to provide thorough coverage. Under severe disease conditions, use the highest rate. Minimum retreatment interval is 7 days.
	Anthraxnose (<i>Collectotrichum acutatum</i>)	8 fl. oz.	Apply MPower Propiconazole in at least 15 gallons of spray per acre beginning at bud break using ground or air equipment in sufficient volume to provide thorough coverage on a 7 -14 day interval.
<p>Almond Restrictions:</p> <ul style="list-style-type: none"> - Do not apply more than 32 fl. oz. per acre per year. - Do not apply more than 0.90 lb. a.i. propiconazole per acre per year. - Do not apply within 60 days of harvest - Do not graze livestock in treated areas or cut treated cover crop for feed 			
Bananas & Plantains	Black Sigatoka (<i>Mycosphaerella fijiensis</i>)	3 fl. oz.	<p>Make applications before disease symptoms appear at the onset of the rainy season. Apply required rate in 10 to 20 gallons of water per acre using ground or air application equipment. Make no more than 2 consecutive applications on a 21 to 25 day schedule before rotating to another labeled product with a different mode of action for at least 2 sprays. If possible, have at least 2 consecutive months "triazole free" during the period of lower disease pressure.</p> <p>Mixing Procedures</p> <p>Oil-in-Water Emulsion: Add the crop oil to the spray tank. Add the emulsifier (0.6 fl. oz. per gal. of oil) and MPower Propiconazole to the spray tank and mix thoroughly for 5 minutes. Add water to the spray tank and mix thoroughly for 15 minutes.</p> <p>Oil Alone: Add crop oil to the spray tank. Add the MPower Propiconazole to the spray tank and mix thoroughly for 5 minutes. Maintain agitation.</p>
<p>Banana and Plantain Restrictions:</p> <ul style="list-style-type: none"> - Do not apply MPower Propiconazole within 100 yards of non-bagged bananas. - Do not apply MPower Propiconazole on bananas or plantains unless they are protected by polyethylene bags. 			

- Do not apply more than 24 fl. oz. of MPower Propiconazole during each growing season (this includes any pre-harvest sprays).
- Do not feed whole bananas and plantains to animals.
- Do not apply more than 0.67 lb. a.i. propiconazole per acre per year.
- A maximum of 8 applications can be made.

Crop	Pests Controlled	Application Rate/Acre	Instructions
Beets, garden	Leaf Spot (<i>Cercospora beticola</i>) Powdery Mildew (<i>Erysiphe polygoni</i>)	3-4 fl. oz.	Begin applications at first sign of disease. Repeat on a 14-day interval. Make no more than 2 consecutive applications before rotating to another registered fungicide with a different mode of action. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Application: MPower Propiconazole is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. MPower Propiconazole may be applied by ground, air or chemigation.

Specific Use Restrictions:

- (1) Do not apply more than 16 fl. oz./A /year of MPower Propiconazole.
- (2) Do not apply within 14 days of harvest.
- (3) Do not apply more than 0.45 lb a.i. propiconazole containing product/A/year.

Crop	Pests Controlled	Application Rate/Acre	Instructions
BERRIES* *Bushberries Bingleberry, Blackberry, Blueberry, Bosenberry, Currants, Dewberry, Elderberry, Gooseberry, Huckleberry, *Caneberries Loganberry, Lowberry, Marionberry, Olallieberry, Red and Black Raspberry, Youngberry Juneberry Lingonberry Salal And cultivars and/or hybrids of these	Mummyberry Disease (<i>Monilinia vaccinocorymbosi</i>)	6 fl. oz.	Make first application beginning at green tip and repeat in 7 to 10 days. If conditions are favorable for disease development, additional applications may need to be made at pink bud and repeated every 7 to 10 days through petal fall.
	Leaf spot and Stem canker (<i>Septoria albopucata</i>) Rust (<i>Pucciniastrum vaccinii</i>)	6 fl. oz.	Apply when conditions favor disease development Repeat applications on a 4 week spray interval.
	Leaf and Cane Spot (<i>Septoria rubif</i>)	6 fl. oz.	Apply as a delayed dormant spray after training in the spring. Repeat this application in the late spring, again at bud break, and again once flowering has begun.
	Powdery Mildew (<i>Microsphaera vaccinii</i>)	6 fl. oz.	Apply at 5 -10% bloom. Repeat this application at full bloom and on a 14 day interval while conditions are favorable for disease development

	Leaf Spot (<i>Septoria</i> spp.)	6 fl. oz.	Make first application any time prior to bloom and again after petal fall. If needed, repeat application just after harvest
--	-----------------------------------	-----------	---

Berry Restrictions:

- Unless directed otherwise for a specific pest, MPower Propiconazole may be applied by either ground in a minimum of 5 gal. per acre or air in a minimum of 15 gal. per acre.
- Do not apply more than 30 fl. oz. per acre per year.
- Do not apply within 30 days of harvest
- Do not apply more than 0.84 lb. a.i. propiconazole per acre per year.

Crop	Pests Controlled	Application Rate/Acre	Instructions
CARROTS	Leaf Blights (<i>Cercospora carotae</i>) (suppression of <i>Alternaria dauci</i>)	4 fl. oz.	Apply when conditions favor disease development. Continue applications on a 7 to 10 day interval using the shorter interval when disease conditions are severe. If desired, a spreader-sticker may be used.
	Powdery Mildew (<i>Erysiphe polygoni</i>)	2 fl. oz. plus chlorothalonil at 0.75 lb. a.i.	Apply with 0.75 lb. a.i. of chlorothalonil per acre. Begin applications when conditions favor disease development. Continue applications on a 7 to 10 day interval.

Carrot Restrictions:

- MPower Propiconazole may be applied by either ground in a minimum of 15 gal. per acre or air in a minimum of 5 gal. per acre
- Do not apply more than 16 fl. oz. per acre per year.
- Do not apply within 14 days of harvest.
- Do not apply more than 0.45 lb. a.i. propiconazole per acre per year.

CELERY AND LEAF PETIOLES SUBGROUP Celery Chinese Celery Cardoon Celtuce Florence Fennel Rhubarb Swiss Chard	Early blight (<i>Cercospora apii</i>) Late blight (<i>Septoria apicola</i>)	4 fl. oz.	Apply on a 7-day schedule either by ground or air. MPower Propiconazole may be tank mixed with an appropriate spreader-sticker. Apply in 10 gals. of water for ground application or 5 gals. of water for aerial application.
--	--	-----------	---

Celery and Leaf Petiole Restrictions:

- Do not apply more than 16 fl. oz. per acre per year.
- Do not apply within 14 days of harvest.
- Do not apply more than 0.45 lb a.i. propiconazole per acre per year.

Crop	Pests Controlled	Application Rate/Acre	Instructions
CEREALS Wheat Barley Rye Triticale Oats	Control of leaf diseases: Rust (<i>Puccinia</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) Leaf blight Glume blotch Tan spot (<i>Pyrenophora riticirepentis</i>) Helminthosporium leaf blight Spot blotch (<i>Bipolaris sorokinina</i>) Barley scald (<i>Rhynchosporium secalis</i>) Barley stripe Net blotch (<i>Pyrenophora teres</i>) Fusarium head blight (suppression only)	4 fl. oz.	Protecting the flag leaf is important for maximizing yield. When applied at 50% to fully emerged, the highest yields are normally obtained. Applications may be made no closer than at 14 day intervals. The use of an oil based adjuvant may improve spray coverage.
	Early Season Suppression of: Tan spot Powdery mildew Glume blotch Leaf Blight (<i>Septoria tritici</i>)	2 - 4 fl. oz.	For early season leaf disease suppression, apply at the specified rate for suppression of listed diseases. Apply in the spring. Make a second application up to Feekes growth stage 10.5 for season long control. Applications may be made no closer than a 14 day interval.
	Foot rot (<i>Pseudocercospora</i> spp.)	4 fl. oz.	Apply the specified rate of MPower Propiconazole per acre plus half rates of other EPA-registered fungicides. Apply at tillering but before elongation has occurred.
Cereal Restrictions: - Applications may be made using ground, air, or chemigation equipment. - Do not apply more than 8 fl. oz. per acre per year. - Do not apply more than 4 fl. oz. per acre per year if forage or hay will be harvested. - Do not apply within 30 days of harvest for forage, 40 days before harvest for grain and straw, and 45 days before harvest for hay. - Do not apply more than 0.22 lb. a.i. propiconazole per acre per year.			

Crop	Pests Controlled	Application Rate/Acre	Instructions
CITRUS (non-bearing) Calamondin Citron Citrus hybrids Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Pummelo Satsuma Mandarin Tangerine Including all cultivars and/or hybrids of these	Greasy Spot	6-8 fl oz	Begin applications in June. Apply at 30 day intervals through August. MPower Propiconazole may be applied by either ground or aerial application in a minimum of 15 gal. per acre.
Citrus Restrictions: - Do not apply more than 24 fl. oz. per acre per year. - Do not apply to citrus that will bear harvestable fruit within 12 months. - Do not apply more than 0.67 lb. a.i. propiconazole per acre per year.			
CORN (FIELD, SEED, AND POPCORN) SWEET CORN	Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>) Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) Also known as Helminthosporium leaf blights (<i>Helminthosporium maydis</i> , <i>H. turcicum</i> , and <i>H. carbonum</i>)	2 - 4 fl. oz.	Apply when disease first appears and continue on a 7 to 14 day schedule. Use the low rate when disease pressure is low. Under heavy pressure or when conditions favor disease development, apply the high rate. Apply by ground, air, or chemigation.
	Rusts (<i>Puccinia</i> spp.) Gray leaf spot (<i>Cercospora zeaemaydis</i>) Eye spot (<i>Aureobasidium zeae</i>)	4 fl. oz.	Apply by ground, air, or chemigation when rust pustules first appear and continue on a 7 to 14 day schedule when conditions favor disease development. For best disease control, early applications at initial disease onset perform better.
Corn Restrictions: For field corn, field corn grown for seed, and popcorn:			

- Do not apply more than 16 fl. oz. per acre per year.
- Do not apply within 30 days of harvest for forage, grain, and stover
- Do not apply more than 8 fl. oz. per acre per year on field corn harvested for forage.
- Do not apply more than 0.45 lb. a.i. propiconazole per acre per year.

For sweet corn:

- Do not apply within 14 days of harvest for ears and 14 days of harvest for forage.

Crop	Pests Controlled	Application Rate/Acre	Instructions
CRANBERRIES (OR, WA, WI only)	Cottonball (<i>Monilinia oxycocci</i>)	4 - 6 fl. oz.	Make the first application at leaf bud break. Make the second application 14 days later. Make the third application at early bloom and repeat again in 14 days. Under severe pressure, use the higher rate for control. Apply by either ground or aerial application equipment in a minimum of 20 gal. per acre.
<p>Cranberry Restrictions:</p> <ul style="list-style-type: none"> - Do not apply more than 24 fl. oz. per acre per year. - Do not apply within 45 days of harvest. - Do not apply more than 0.67 lb. a.i. propiconazole per acre per year. 			
FILBERTS (Hazelnuts)	Eastern Filbert Blight (<i>Anisogramma anomala</i>)	5 - 8 fl. oz.	Begin applications when green leaf tissue becomes visible and continue at 14 to 21 day intervals. Under severe disease conditions, use the higher rate and shorter interval. On certain varieties, MPower Propiconazole applications may cause smaller and/or greener leaves. Yields of filberts displaying these characteristics have not been reduced due to propiconazole treatments. Apply by either ground or aerial application in a minimum of 15 gal. per acre
<p>Filbert Restrictions:</p> <ul style="list-style-type: none"> - Do not apply more than 32 fl. oz. per acre per year. - Do not apply more than 0.90 lb. a.i. propiconazole per acre per year. - Do not apply within 60 days of harvest. - Do not graze livestock in treated areas or cut treated crop for feed. 			
GRASSES GROWN FOR SEED (NEBRASKA, OREGON, WASHINGTON, IDAHO, AND MINNESOTA ONLY)	Rusts (<i>Puccinia</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) Selenophoma stem eyespot (<i>Selenophoma</i>) Ergot Suppression	4 - 8 fl. oz.	Apply by ground, by air in a minimum of 10 gals. of water per acre, or through irrigation equipment. Apply when powdery mildew and Selenophoma infections or rust pustules are noticeable and increasing in number in late spring or early summer. Repeat at 14 to 21 day intervals. To maximize control under severe rust pressure, use the higher rate of 8 fl. oz. per acre and make applications at 14 day intervals until the seed is mature.

			Make the last application at least 20 days before seed matures. For bluegrass, it is important to begin applications early in the growing season.
--	--	--	---

Grasses Grown for Seed Restrictions:

- Do not apply more than 32 fl. oz. per acre per growing cycle.
- Do not feed hay cut within 20 days of the last application.
- Do not graze treated areas within 140 days of the last application.
- Do not apply more than 0.90 lb. a.i. propiconazole per acre per year.

Crop	Pests Controlled	Application Rate/Acre	Instructions
MINT Peppermint Spearmint	Rust <i>(Puccinia menthae)</i>	4 fl.oz.	Apply in a minimum of 20 gals, of water per acre using ground application. Begin applications when plants are 2 to 4 inches high or when conditions become favorable for disease development. Make a second application 14 days after the first application.

Mint Restrictions:

- Do not apply within 30 days of harvest.
- Do not exceed 8 fl. oz. per acre per year.
- Do not apply more than 0.22 lb. a.i. propiconazole per acre per year.

ONIONS (dry bulb) GARLIC SHALLOTS (dry bulb) ONIONS, GREEN Green Shallots Green Eschalots Japanese bunching onions Leeks Spring Onions Scallions And or cultivars or hybrids of these	Purple Blotch <i>(Altemari pom)</i> Suppression of Botrytis leaf blight <i>(Botrytis squamosa)</i>	4 - 8 fl. oz. 2 - 4 fl. oz. plus tank mix partner	Apply by ground in a minimum of 15 gals. of water per acre, or by air in a minimum of 5 gals. of water per acre. Begin applications when conditions favor disease development and continue on a 7 to 10 day interval. Use the higher rate and shorter interval when disease conditions are severe. In tank mix, apply specified rate in combination with another fungicide for control of Botrytis leaf blight or purple blotch. Begin applications when conditions favor disease development and continue on a 7 to 10 day interval or according to the tank mix partner's label. Use higher rates when disease conditions are severe. To achieve optimum control, use a wetting agent or a spreader/sticker.
--	---	--	--

Onion, Garlic, Shallot, and Green Onion Restrictions:

- Do not apply more than 16 fl. oz. per acre per year.
- Do not apply within 14 days of harvest on dry bulb onions.
- May be applied on the day of harvest for green onion types
- Do not apply more than 0.45 lb. a.i. propiconazole per acre per year.

Crop	Pests Controlled	Application Rate/Acre	Instructions
Parsley, fresh and dried leaves Cilantro (Coriander), leaves	Leaf Spot <i>(Cercospora spp.)</i> Leaf Spot <i>(Alternaria spp.)</i> Powdery Mildew <i>(Erysiphe spp.)</i>	3-4 fl. oz.	Begin applications at first sign of disease. Repeat on a 14-day interval. Make no more than 2 consecutive applications before rotating to another registered fungicide with a different mode of action. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.

Application: MPower Propiconazole is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. MPower Propiconazole may be applied by ground, air or chemigation.

- Specific Use Restrictions:**
- 1) Do not apply more than 16 fl. oz./A/year of MPower Propiconazole.
 - 2) Do not apply within 14 days of harvest.
 - 3) Do not apply more than 0.45 lb a.i. propiconazole containing products/A/year.

Crop	Pests Controlled	Application Rate/Acre	Instructions
PEANUTS	Late leaf spot <i>(Cercosporidium)</i> Early leaf spot <i>(Cercospora arachidola)</i> Rust <i>(Puccinia arachidis)</i> Web Blotch <i>(Phoma arachidicola)</i>	2.5 - 4 fl. oz.	Use 2.5 - 4 fl. oz. on Early leaf spot Use 4 fl. oz. on all other listed diseases. Apply MPower Propiconazole alone using ground, aerial, or chemigation equipment beginning applications 35 to 40 days after planting or at the first appearance of disease. Continue applications on a 10 to 14 day schedule. Under heavy disease pressure, use higher application rates MPower Propiconazole also may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.
	Southern Stem Rot <i>(Sclerotium rolfsii)</i>	See Instructions section for appropriate rate	Apply according to one of the following schedules: A. Apply 4 fl. oz. per acre to the crown and pegging zones of the plant using chemigation or directed ground application. Begin applications 45 days after planting or at the first appearance of disease, and repeat on a 14 day schedule. B. Apply 8 fl. oz. per acre to the crown and pegging zones of the plant using chemigation or directed ground application.

		<p>Make 2 applications; the first at pegging (approximately 60 days after planting) or at the first appearance of disease, and the second application 3 to 4 weeks later.</p> <p>Irrigation: When applying in irrigation water for Southern Stem Rot Control, use a minimum of 0.25 to 0.5 inch of irrigation water per acre. Use enough water so that the fungicide penetrates the peanut canopy and reaches the crown of the plant where <i>Sclerotium rolfsii</i> is most active.</p> <p>When using this product via irrigation or directed ground application, additional methods should be used for leaf spot control.</p>
--	--	--

Peanut Restrictions:

- Do not apply more than 16 fl. oz. per acre per year.
- Do not feed hay from treated fields to livestock if the high rate (8 fl. oz.) was used.
- Do not apply within 14 days of harvest when using no more than 4 fl. oz. per acre and within 21 days of harvest using 8 fl. oz. per acre.
- Do not apply more than 0.45 lb. a.i. propiconazole per acre per year.

Crop	Pests Controlled	Application Rate/Acre	Instructions
PECANS	Pecan Scab <i>(Cladosporium caryigenum)</i> Downy Spot <i>(Mycosphaerella caryigena)</i> Liver Spot <i>(Gnomonia caryae pv pecanae)</i> Vein Spot <i>(Gnomonia nerviseda)</i> Zonate Leaf Spot <i>(Cirstulariella moricola)</i> Powdery Mildew <i>(Microsphaera penicillata)</i>	4 - 8 fl. oz.	<p>Pecan scab: Apply 4 - 8 fl. oz. per acre on a 14 day schedule during bud break and pre-pollination sprays. Apply 6-8 fl. oz. per acre during nut formation and coyer sprays. Use higher rates when disease pressure is heavier.</p> <p>Other listed foliar diseases: Apply 4 fl. oz. per acre with other registered pecan products labeled for these mid to later season foliar diseases. Observe all directions, precautions, and limitations for the other products. MPower Propiconazole may be applied by either ground or by aerial application in a minimum of 20 gal. per acre. Propiconazole may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are</p>

			making the pesticide application. To determine whether your county has a bulletin, consult http://www.epa.gov/espp/usa-map.htm . Bulletins may also be available from local pesticide dealers, extension offices, or state pesticide agencies
--	--	--	---

Pecan Restrictions:

- Do not apply more than 32 fl. oz. per acre per year.
- Do not apply after shuck split.
- Do not apply more than 0.9 lb. a.i. of propiconazole per acre per year.
- Do not graze livestock in treated areas or cut treated cover crop for feed
- Do not apply within 30 days of harvest.

PINEAPPLE (HAWAII ONLY)	butt rot disease of pineapple (<i>Ceratocystis paradoxa</i>)	0.75 fl. oz. (22 ml) per 100 gals. of water (117,000)	Treatments can be made in either a cold or hot water dip. Cold Water Dip -Immerse crowns to give thorough wetting, remove, and allow to drain. Hot Water Dip -Maintain water temperature at 125°F (52°C). Soak crowns for 20 to 30 minutes, remove, and allow to drain.
-----------------------------------	---	---	---

Pineapple Restrictions:

- Do not use treated crowns for food or feed
- Do not graze while plant is growing
- Do not graze tops until fruit is harvested.
- Dispose of used dip solution according to local, state, and federal regulations.

Crop	Pests Controlled	Application Rate/Acre	Instructions
PISTACHIOS	Botryosphaeria Panicle and Shoot Blight (<i>Botryosphaeria dothidea</i>)	5 - 8 fl oz	Begin applications when green leaf tissue becomes visible and continue on a 14 to 21 day interval. Under severe disease conditions, use the higher rate and the shorter interval. Under certain conditions MPower Propiconazole applications may cause smaller and/or greener leaves. Yields of pistachios displaying these characteristics have not been reduced due to MPower Propiconazole treatments. Apply by either ground or by aerial application in a minimum of 15 gal. per acre.

Pistachio Restrictions:

- Do not apply more than 32 fl. oz. per acre.
- Do not apply within 60 days of harvest
- Do not apply more than 0.90 lb. a.i. propiconazole per acre per year.
- Do not graze livestock in treated areas or cut treated cover crop for feed.

RICE	Sheath blight <i>(Rhizoctonia solani)</i> Brown leaf spot <i>(Helminthosporium oryzae)</i> Narrow brown leaf spot and brown blotch <i>(Cercospora oryzae)</i> Leaf smut <i>(Entyloma oryzae)</i> Sheath spot <i>(Rhizoctonia oryzae)</i> Kernal smut <i>(Tilletia barclayana)</i> Aggregate sheath spot <i>(Rhizoctonia oryzaesativa)</i> Black sheath rot <i>(Gaeumannomyces graminis)</i> Stem rot suppression <i>(Sclerotium oryzae)</i> False smut suppression <i>(Ustilaginoida virens)</i>	See Instructions section for appropriate rate	The timing of application will depend on disease severity, disease complex, and rice variety and growth stage. Apply MPower Propiconazole at specified rates on either of the following schedules as an aerial spray in 5 to 10 gals. of water per acre: A. 6 fl. oz. per acre at first internode elongation (up to 2 inch panicle) and repeat at swollen boot. Make the second application 10 to 14 days after the first application, but before the boot splits and head emerges. MPower Propiconazole provides best control of sheath blight when the first application is applied at disease appearance in the field. Make the first application when 5% or fewer of the tillers are infected. B. 10 fl. oz. per acre at first internode elongation (up to 2 inch panicle). Use the 10 oz. rate if greater than 10% of the tillers are infected with sheath blight. If disease reappears, use another registered fungicide for the second application. C. Apply 6 fl. oz. per acre in a tank mix with Quadris® or other fungicides for control of diseases of rice.
-------------	---	--	---

Crop	Pests Controlled	Application Rate/Acre	Instructions
Wild Rice (MN only)	Helminthosporium leaf blight and brown spot <i>(Bipolaris spp.)</i>	6 - 8 fl. oz.	Apply specified rate of MPower Propiconazole per acre at both booting and heading, or make a single application of 8 fl. oz. per acre at booting. Make application using aerial application equipment. The minimum application interval is 10 days. For Rice and Wild Rice Uses: Propiconazole may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product you must follow the measures contained in the County Bulletin for the county in which you are making the pesticide application. To determine whether your county has a bulletin, consult http://www.epa.gov/espp/usa-map.htm . Bulletins may

	also be available from local pesticide dealers, extension offices, or state pesticide agencies.		
Rice and Wild Rice Restrictions:			
<ul style="list-style-type: none"> - Do not make applications using ground or chemigation equipment. - Only aerial application is allowed. - Do not apply to stubble or ratoon crop rice. - Do not use in rice fields where commercial farming of crayfish will be practiced. - Do not drain water from treated rice fields into ponds used for commercial fish farming. - Do not use water drained from treated fields to irrigate other crops. - Do not apply more than 12 fl. oz. per acre per year. - Do not apply within 45 days of harvest. - Do not apply more than 0.34 lb. a.i. propiconazole per acre per year. - Do not apply within 10 days of harvest for wild rice. 			
Sorghum	Ergot (<i>Claviceps sorghi</i>)	3-4 fl. oz.	Make first application at or just prior to flowering. Repeat on a 5 to 7 day interval. Apply up to four times. Make application using aerial application equipment in a minimum of 10 gals. of spray per acre or by ground in a minimum of 15 gals. of spray per acre.
Sorghum Restrictions:			
<ul style="list-style-type: none"> - Do not apply more than 16 fl. oz. per acre per year. - Do not apply within 30 days of harvest for forage. - Do not apply within 21 days of harvest for grain and stover - Do not graze livestock or cut for green chop or silage within 30 days of application. - Do not apply more than 8 fl. oz. per acre per year on sorghum harvested for forage. - Do not apply more than 0.45 lb. a.i. propiconazole per acre per year. 			
Crop	Pests Controlled	Application Rate/Acre	Instructions
Soybeans	Aerial Web Blight (<i>Rhizoctonia solani</i>) Anthracnose (<i>Colletotrichum truncatum</i>) Brown Spot (<i>Septoria glycines</i>) Frogeye Leaf Spot (<i>Cercospora sojina</i>) Soybean Rust (<i>Phakopsora pachyrhizi</i>)	4 - 6 fl. oz.	Applications may be made using ground or aerial application equipment. Use dilution rates found in the APPLICATION INSTRUCTIONS section of this label. When applying by air, adding an oil-based additive improves coverage and penetration. Apply 5 - 6 fl. oz. at the first appearance of Aerial web blight and repeat the application 14 to 21 days later. Under severe conditions, use the higher rate and shorter interval. For control of other foliar diseases, apply 6 fl. oz. at growth stage R3 (early pod set) when pods are 1/8 to 1/4 inch long and 21 days later at growth stage R5 (pod fill). Apply 4 - 6 fl. oz. at first indication that soybean rust is in the area. For best control, preventative applications work best. Repeat on a 14 to 21 day interval using the higher

			rate and shorter interval when disease is present in field and incidence is less than 2% (2 plants in 100 infected). If incidence is greater than this or if disease is in mid canopy, control will not be acceptable. Scouting for rust and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications. On certain varieties, MPower Propiconazole applications may cause crinkled or smaller greener leaves. Yields of dry beans displaying these characteristics have not been reduced due to propiconazole treatments
--	--	--	--

Soybean Restrictions:

- Do not apply more than 12 fl. oz. per acre per year.
- Applications may be made up to growth stage R6.
- Do not apply more than 0.34 lb. a.i. propiconazole per acre per year.
- 30 day PHI.
- Apply in a minimum of 5 and 15 gal/A using aerial and ground equipment, respectively.
- Do not graze or feed soybean, forage, or hay.

Crop	Pests Controlled	Application Rate/Acre	Instructions
Stone Fruit: Apricots, Cherries (Sweet and Tart), Nectarines, Peaches, Plums, Plumcots, Prunes And Cultivars and/or Hybrids of These	Brown Rot Blossom Blight (<i>Monilinia</i> spp.)	4 fl. oz.	Apply by ground or air in a minimum of 15 gal. per acre at early bloom stage. Stone fruit diseases are most effectively controlled by ground applications. If disease pressure is low, a second application may be made as needed up through petal fall. Make a second application if disease pressure is high or for susceptible varieties at 75 - 100% bloom. If blossoming is prolonged or conditions favorable for disease persist, make a third application at petal fall.
	Powdery Mildew (<i>Podosphaera</i> spp.) Cherry Leafspot (<i>Blumeriella jaapii</i>) Rust (<i>Tranzschelia discolor</i>)	4 fl. oz.	Follow the brown rot blossom blight schedule above applying by ground or air in a minimum of 15 gal. per acre. Stone fruit diseases are most effectively controlled by ground applications. Make up to 2 additional applications on a 10 to 14 day interval from the end of petal fall to harvest.
	Fruit Brown Rot (<i>Monilinia</i> spp.)	4 fl. oz.	Apply by ground or air in a minimum of 15 gal. per acre as needed with a maximum of 2 sprays during the preharvest period up to the day of harvest (0 day PHI). Stone fruit

			diseases are most effectively controlled by ground applications. If high inoculum and severe disease conditions persist, apply another registered fungicide after the two MPower Propiconazole applications.
--	--	--	--

Stone Fruit Restrictions:

- MPower Propiconazole may be applied on the day of harvest.
- Do not apply more than 0.225 lb. propiconazole per acre per year.
- Do not apply more than 20 fl. oz. per acre per year.
- Applications made during bloom to Stanley plums have occasionally caused fruit to be less oval in shape and smaller in size at harvest. To avoid this, do not apply to Stanley plums earlier than 21 days before harvest.
- Do not apply within 10 days of harvest.

Strawberries	Anthracnose (<i>Colletotrichum acutatum</i>) Leaf Spot (<i>Cercospora fragariae</i>) Powdery Mildew (<i>Sphaerotheca macularis</i>) Leaf Rust (<i>Phragmidium potentillae</i>)	4 fl. oz.	Begin applications when disease levels are no more than 5%. Apply up to 4 times on a 7-day interval. Make no more than 2 consecutive applications before rotating to another registered fungicide with a different mode of action. This product may be applied by either ground in a minimum of 20 gal. per acre or aerial in a minimum of 15 gal. per acre.
---------------------	---	-----------	--

Strawberry Restrictions:

- Do not apply more than 16 fl. oz. per acre per year.
- May be applied on the day of harvest.
- Do not apply more than 0.45 lb. a.i. propiconazole per acre per year.

Crop	Pests Controlled	Application Rate/Acre	Instructions
Sugar Beets	Leaf Spot (<i>Cercospora beticola</i>) Powdery Mildew (<i>Erysiphe polygoni</i>)	4 fl. oz.	Begin applications are first sign of disease and repeat at 10 to 14 day intervals. Make no more than 2 consecutive applications before rotating to another registered fungicide with a different mode of action. If disease levels continue to increase, immediately switch to a fungicide with a different mode of action. This product may be applied by air, ground, or chemigation equipment. Use dilution rates found in the APPLICATION INSTRUCTIONS section of this label.

Sugar Beet Restrictions:

- Do not apply more than 12 fl. oz. per acre per year.
- Do not apply within 21 days of harvest
- Do not apply more than 0.34 lb. a.i. propiconazole per acre per year.

Sugarcane	Pineapple disease (<i>Ceralocystis paradoxa</i>)	0.75 fl. oz. (22 ml) per 100 gals. of water (1:17,000)	For this use only in Hawaii. Apply this product to cut seed pieces. Treatments can be applied in either a cold or hot water dip. Cold Water Dip -Immerse seed pieces to give thorough wetting, remove, and allow to drain. Hot Water Dip -Maintain water temperature at 125°F (52°C). Soak the seed pieces for 20 to 30 minutes, remove, and allow to drain. Conveyor Belt Treatment -Treat seed pieces with MPower Propiconazole /water solution using in-line directed spray sufficient to wet cut ends.
------------------	---	--	--

Sugar Cane Restrictions:

- Do not use treated seed pieces for food or feed purposes.
- Dispose of spent dip solution according to state and federal regulations.

Crop	Pests Controlled	Application Rate/Acre	Instructions
<p>TREE NUTS Almond (see specific directions in ALMOND Section) Beechnut Brazil Nut Butternut Cashew Chestnut Chinquapin Filbert (see specific directions in FILBERT section) Hickory Macadamia Pecan (see specific directions in PECAN section) Walnut Pistachios (see specific directions in PISTACHIO section)</p>	<p>Foliar Diseases</p>	<p>4 - 8 fl. oz.</p>	<p>Apply at first sign of disease. Repeat on a 7 to 14 days interval. May be applied by either ground or aerial application in a minimum of 15 gal. per acre. Tree nut diseases are most effectively controlled by ground applications.</p>
<p>Tree Nut Restrictions:</p> <ul style="list-style-type: none"> - Do not apply more than 32 fl. oz. per acre per year. - Do not apply within 60 days of harvest except for pecan (see specific directions in PECAN section of this label). - Do not apply more than 0.90 lb. a.i. propiconazole per acre per year. - Do not graze livestock in treated areas or cut treated cover crop for feed 			

POST HARVEST USE

Crop	Pests Controlled	Use Rate fl. oz. product/100 gals.	Instructions
Pineapple	Ceratocytis Butt Rot (<i>C. paradoxa</i>)	3-4	Apply 3 to 4 fluid ounces in 100 gallons of water or wax water emulsion after harvest. Fruit should be dipped or sprayed for thorough coverage and allowed to drain. Limit dipping time to no more than 3 minutes. Dip solution should be replaced with fresh dip solution after 200,000 lbs of fruit have been treated. Fruit discarded from fresh fruit packing operations may be used for processing Cannery wastes may be fed

Specific Use Restrictions:

- 1) Do not apply more than 4 fl. oz. as a post harvest treatment.

Fl. Oz. MPower Propiconazole Per Acre	lb. A.I. per Acre	Acres Treated Per 1 Gallon of MPower Propiconazole
2	0.056	64.0
4	0.1125	32.0
6	0.169	21.3
8	0.225	16.0
10	0.28	12.8
12	0.34	10.7
16	0.45	8.0
20	0.56	6.4
24	0.67	5.3
30	0.84	4.3
32	0.90	4.0

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

INFORMATION FOR TURFGRASS AND ORNAMENTAL USES

MPower Propiconazole is a systematic fungicide for use on turfgrasses for the control of dollar spot (*Sclerotinia homeocarpa*), brown patch (*Rhizoctonia solani*), anthracnose (*Colletotrichum graminicola*), red thread (*Laetisaria fuciformis*), pink patch (*Limonomyces roseipellis*), rust (*Puccinia graminis*), powdery mildew (*Erysiphe graminis*), stripe smut (*Ustilago striformis* and *Urocystis agropyri*), summer patch (*Magnaporthe poae*), necrotic ring spot (*Leptosphaeria korrae*), spring dead spot (*Leptosphaeria korrae*, *Leptosphaeria narmari*, *Ophiosphaerella herpoticha*, *Gaeumannomyces graminis*), take-all patch (*Gaeumannomyces graminis*), leafspot (*Bipolaris* spp., *Drechslera* spp.), gray leafspot (*Pyricularia grisea*), pink snowmold (*Microdochium nivale*), Fusarium patch (*Fusarium nivale*), gray snowmold (*Typhula* spp.), yellow patch (*Rhizoctonia cerealis*), and zoysia patch (*Rhizoctonia solani*).

MPower Propiconazole also controls numerous diseases on ornamentals and other landscape and nursery plantings. It controls powdery mildews, rusts, leafspots, scabs, and blights. Refer to the appropriate section for specified diseases and plants.

Restrictions: For turfgrass and ornamental uses, do not apply this product through any type of irrigation system. Do not use MPower Propiconazole in greenhouses or as a tree injection. Do not apply more than 5.8 fl. oz. per 1,000 sq. ft. of MPower Propiconazole per calendar year.

MIXING INSTRUCTIONS

Fill the spray tank $\frac{1}{2}$ to $\frac{3}{4}$ full with water. Add the proper amount of MPower Propiconazole and then add the rest of the water. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

If MPower Propiconazole is tank mixed with other products, use the following sequence:

1. Always check the compatibility of the tank mix using a jar test with proportionate amounts of MPower Propiconazole, other chemicals to be used, and the water, before mixing in the spray tank.
2. Provide sufficient jet or mechanical agitation during filling and application to keep the tank mix uniformly suspended.
3. Fill tank at least $\frac{1}{2}$ full of clean water.
4. Add wettable powders to the tank first, allowing them to completely suspend in the tank before proceeding. This process can be hastened by premixing the product in water before adding to the tank.
5. Add flowables or suspensions next.
6. Add MPower Propiconazole next.
7. Add emulsifiable concentrates last.
8. Do not leave tank mix combinations in the spray tank for prolonged periods without agitation. Mix and apply the same day.

TANK MIXES

For broader spectrum control, MPower Propiconazole can be tank mixed with other fungicides. MPower Propiconazole is also compatible with numerous herbicides and insecticides. Check compatibility before tank mixing. Add Unite® (3 pts. per 100 gals.) to tank mixes which are incompatible. Follow the directions under MIXING INSTRUCTIONS section of this label for tank mixes. Observe all directions, precautions, and limitations on labeling of all products used in tank mixes. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

TURFGRASS AND DICHONDRA DISEASE CONTROL

- **USE MPOWER PROPICONAZOLE IN A PREVENTATIVE DISEASE CONTROL PROGRAM**
- Apply sufficient water to ensure thorough coverage.
- Apply after mowing or allow sprayed area to completely dry before mowing.
- For control of foliar diseases, allow sprayed area to completely dry before irrigation.
- For control of soil-borne diseases, MPower Propiconazole can be watered in after application.
- Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
- For optimum turf quality and disease control, use MPower Propiconazole in conjunction with turf management practices that promote good plant health and optimum disease control.
- Evaluate spray additives prior to use. Label directions are based on data obtained with no additives.
- Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.
- Bermudagrass can be sensitive to MPower Propiconazole. Do not exceed 1.44 fl. oz. per 1,000 sq. ft. every 30 days on any variety of bermudagrass. In Florida, do not apply MPower Propiconazole to bermudagrass golf course greens when temperatures exceed 90F.

RESTRICTIONS:

- Do not apply more than 5.8 fl. oz. per 1,000 sq. ft. per calendar year nor apply more than 1.79 lb a.i. per acre per application.
- Do not graze animals on treated areas. Do not feed clippings from treated areas to livestock or poultry.

Turfgrass – Specific Diseases, Rates, and Application Timing

Disease	Fl. Oz. Per 1,000 Sq. Ft.	Fl. Oz. Per Acre	Application Interval/Timing	Instructions
Dollar Spot (<i>Sclerotinia homeocarpa</i>)	0.18	8.0	7 days	Apply when conditions are favorable for disease development.
	0.18	8.0	14 days	Tank mix with low label rate of one of the following fungicides: Daconil Weatherstik®, Daconil Ultrex®
	0.37	16	21-28 days	Tank mix with low label rate of one of the following fungicides: Daconil Weatherstik®, Daconil Ultrex®, Chipco 26019
	0.37-0.73	16-32	14-28 days	If using the 0.37-0.73 fl. oz. per 1,000 sq. ft. rate without tank mixing, make no more than 3 consecutive applications for dollar spot control before rotating to an alternate EPA registered fungicide having a different mode of action
Antracnose (<i>Colletotrichum graninicola</i>)	0.37-0.73	16-32	14-28 days	Apply when conditions are favorable for disease development. When disease pressure is high, use higher rates of MPower Propiconazole and shorter intervals. For broad spectrum control, tank mix with a registered contact fungicide at the label rate. If disease is present, mix 0.73 fl. oz. of MPower Propiconazole per 1,000 sq. ft. with the label rate of the above mentioned fungicide.
Brown Patch (<i>Rhizoctonia solani</i>)	0.37-0.73	16-32	14-21 days	Begin applications in May or June before disease is present. Tank mix with a registered contact fungicide labeled for Brown Patch control at the label rate. Under conditions of high temperatures and high humidity, use the higher rates of MPower Propiconazole and shorter intervals.
Powdery Mildew (<i>Erysiphe graminis</i>), Rust (<i>Puccinia graminis</i>)	0.37-0.73	16-32	14-28 days	Apply when conditions are favorable for disease development. If disease is present, use 0.73 fl. oz. of MPower Propiconazole per 1,000 sq. ft.
Red Thread (<i>Laetisaria fuciformis</i>) Pink Patch (<i>Limonomyces roseipellis</i>)	0.37	32	14-21 days	Apply when conditions are favorable for disease development.

Disease	Fl. Oz. Per 1,000 Sq. Ft.	Fl. Oz. Per Acre	Application Interval/Timing	Instructions
Stripe Smut (<i>Ustilago striiformis</i>) (<i>Urocystis agropyri</i>)	0.37- 0.73	16-32	Fall or Spring	Apply once in the fall after grass becomes dormant or in the early spring before grass starts to grow.
Gray Leafspot (<i>Pyricularia grisea</i>)	0.37- 0.73	16-32	14 days	Apply when conditions are favorable for disease development. If using the .037 fl. oz. per 1,000 sq. ft. rate, tank mix with a registered contact fungicide at the label rate
Melting out, Leaf Spot (<i>Bipolaris spp.</i>) (<i>Drechslera spp.</i>)	0.37- 0.73	16-63	14 days	Under light to moderate pressure, apply MPower Propiconazole to reduce the severity of leaf spot and melting out. For broad spectrum disease control tank mix 0.37 fl. oz. of MPower Propiconazole rate with a registered contact fungicide at the label rate Tank mix the 0.37-0.73 fl. oz. per 1,000 sq. ft. MPower Propiconazole rate with a registered contact fungicide at the label rate
Summer Patch, Poa Patch (<i>Magnaporthe poae</i>)	0.73 1.45	32 63	14 days 28 days	Apply MPower Propiconazole beginning in April. Use the 1.45 fl. oz. per 1,000 sq. ft. rate on a 28 day schedule and the 0.73 fl. oz. per 1,000 sq. ft. rate on a 14 day schedule.
Take-All Patch (<i>Gaeumannomyces graminis</i>)	0.73- 1.45	32-63	Spring and Fall	Apply MPower Propiconazole to reduce the severity of take-all patch. Make one to 2 fall applications in September and October or when night temperatures drop to 55°F, and 1 to 2 spring applications in April and May, depending on local recommendations.
Spring Dead Spot (<i>Leptosphaeria korrae</i> , <i>Leptosphaeria narmari</i> , <i>Ophiosphaerella herpotricha</i> , <i>Gaeumannomyces graminis</i>)	1.45	63	30 days	Make 1 to 3 applications. For 1 application, apply in September or October. For multiple applications, begin sprays in August.
Necrotic Ring Spot (<i>Leptosphaeria korrae</i>)	1.45	63	Fall or Spring	Apply in the fall and/or the early spring depending on local recommendations.
Snowmold, Gray (<i>Typhula spp.</i>) Pink (<i>Microdochium nivale</i>)	0.73- 1.45	32-63	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow. For optimum disease control, the 0.73-1.45 fl. oz. MPower Propiconazole rate should be tank

Disease	Fl. Oz. Per 1,000 Sq. Ft.	Fl. Oz. Per Acre	Application Interval/Timing	Instructions
				mixed with either PCNB or chlorothalonil at label rates
Fusarium Patch (<i>Fusarium nivale</i>)	0.73-1.45	32-63	Fall-Early Spring	Apply when conditions are favorable for disease development.
Yellow Patch (<i>Rhizoctonia cerealis</i>)	1.10-1.45	48-63	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow. If using a 1.10 fl. oz. per 1,000 sq. ft. rate, tank mix with a registered contact fungicide at the label rate
Zoysia Patch, large patch of zoysia (<i>Rhizoctonia solani</i>)	1.10-1.45	48-63	Early Fall	Make one application in the early fall (mid-September to mid-October) prior to development of disease symptoms. Consult local turfgrass extension experts to determine optimum application timing for your area.

DICHONDRA – Specific Diseases, Rates, and Application Timing

Disease	Fl. Oz. Per 1,000 Sq. Ft.	Fl. Oz. Per Acre	Application Interval/Timing	Instructions
Dichondra Rust (<i>Puccinia dichondrae</i>)	0.73	32	14-21 days	Apply when conditions are favorable for disease development.

Establishment of Cool Season Turfgrass

MPower Propiconazole provides control of many diseases of turf, and its primary use is as a fungicide for use against the diseases listed on this label. As an additional benefit, MPower Propiconazole will improve the establishment rate when it is applied to cool season grass seedlings or sod.

New Seedlings: Apply 0.35 fl. oz. per 1,000 sq. ft. at the 2 to 3-leaf stage of growth for faster root development and top growth. **Sod:** Apply 0.35 fl. oz. per 1,000 sq. ft. 2 to 6 weeks before cutting for increased sod knitting and faster establishment after laying.

DISEASE CONTROL IN NURSERIES (FIELD) AND LANDSCAPE PLANTINGS

- Use MPower Propiconazole in a preventative disease control program.
- To determine the use directions for controlling a disease on an ornamental plant species, select the plant species in Table 1. The number in parenthesis following the plant species refers you to the disease(s) controlled in Table 2. Find the disease in Table 2. The letter in brackets following the disease refers you to the application regime in Table 3.
- Allow spray to dry before overhead irrigation is applied.
- Optimum benefit of MPower Propiconazole is obtained when used in conjunction with sound disease management practices.

APPLICATION DIRECTIONS

MPower Propiconazole may be used at rates of 0.75-8.7 fl. oz. per 100 gallons of water for control of diseases of ornamental plant species (see Tables 1, 2, and 3).

For outdoor uses, you can apply up to 2.0 gallons of MPower Propiconazole per acre per crop per calendar year.

For disease control in landscapes, apply 2.2-3.0 fl. oz. per 100 gallons of water every 21 days. For best control, begin MPower Propiconazole applications before disease development.

Plant tolerances to MPower Propiconazole have been found to be acceptable for the specific genera and species of plants listed under the Directions for Use. Other plant species may be sensitive to MPower Propiconazole and diseases other than those listed may not be controlled. Before using MPower Propiconazole on plants or for diseases that are not listed in the Directions for Use, test MPower Propiconazole on a small-scale basis first. Do not apply MPower Propiconazole to African violets, begonias, Boston fern, or geraniums. Apply the recommended rates for a particular type of disease, i.e., rust, powdery mildew, etc., and evaluate for phytotoxicity and disease control prior to widespread use.

Table 1. Ornamentals - Plant Species

Numbers in parenthesis refer to diseases controlled. See Table 2.

Herbaceous Ornamental	Woody Ornamental	Non-bearing Fruits and Nuts (Nurseries and Landscape Plantings)
Calendula (4a)	Amelanchier (4d)	Apple (3q, 4d, 5a)
Carnation (5f)	Ash (4c)	Bartlett pear (3q, 4c, 5a)
Chrysanthemum (2a)	Azalea (2c, 4b)	Cherry (2b, 3d)
Delphinium (4a)	Bayberry(3n)	Citrus (3m)
English Ivy (3e)	Camellia (3e)	Nectarine (2b)
Gomphrena (3a)	Cotoneaster (3i)	Peach (2b)
Impatiens (3a, 3b, 4a)	Crabapple(3c,3q,4c,5a)	Pecan (3b, 3c, 3f, 3l, 3n, 4e)
Iris (5d)	Crape myrtle (4a)	Plum (2b)
Marigold (3a)	Dogwood (3h, 4c)	Walnut (3j)
Monarda (4c)	Douglas fir (5b)	
Phlox (4c)	Elm (4c)	
Snapdragon (5d)	Euonymus (3e, 4c)	
Sweet William	Hawthorn (5a)	
(<i>Dianthus barbatus</i>) (3k)	Holly (3r)	
Zinnia (4c)	Juniper (1a)	
	Lilac (4c)	
	Linden (3e, 3b, 4b)	
	Magnolia (3e, 4b)	
	Maple (3e, 4f)	
	Oaks(3p)	
	Pines (1b, 1c)	
	Poplars (5b)	
	Pyracantha (3o)	
	Red Tip Photinia (3i)	
	Rhaphiolepis (3e, 3i)	
	Rhododendron (2c, 3n)	
	Roses (3g, 4e, 5c)	
	(Outdoor use only)	
	Shasta fir (5e)	
	Sweetgum (3b, 3c, 3n)	
	Sycamore (3e)	
	Tulip tree (3e, 4a)	
	Wax myrtle (3n)	

Table 2. Diseases

Letters in brackets refer to application regimes. Refer to Table 3.

1. Conifer Blights
 - a. *Phomopsis juniperovora* (Phomopsis Blight) [B]
 - b. *Sirococcus strobolinus* (Tip Blight) (D)
 - c. *Sphaeropsis sapinea* (Diplodia Tip Blight) (B)
2. Flower Blight
 - a. *Ascochyta chrysanthemi* (Ray Blight) [C]
 - b. *Monilinia* spp. [A]
 - c. *Ovulinia* spp. [B]

3. Leaf Blights/Spots
 - a. *Alternaria* spp. [B]
 - b. *Cercospora* spp. (Brown Leaf Spot) [C]
 - c. *Cladosporium* spp. (Scab) [C]
 - d. *Coccomyces hiemalis* [A]
 - e. *Colletotrichum* spp. [B]
 - f. *Cristulariella* spp. (Zonate leafspot) [C]
 - g. *Diplocarpon rosae* (Blackspot) [B]
 - h. *Discula* spp. (Anthracnose) [A]
 - i. *Fabraea maculate* (syn. *Entomosporium maculate*) [B]
 - j. *Gnomonia leptostyla* (Anthracnose) [C]
 - k. *Heterosporium echinulatum* [B]
 - l. *Mycosphaerella caryigena* (Downy Spot) [C]
 - m. *Mycosphaerella fructicola* (Greasy Spot) [E]
 - n. *Septoria* spp. (Leaf Scorch) [C]
 - o. *Spilocaea pyracanthae* [B]
 - p. *Tubakia dryina* [D]
 - q. *Venturia inaequalis* (Scab) [A]
 - r. *Rhizoctonia* web blight [B]

4. Powdery
Mildew

- a. *Erysiphe* spp. [B]
- b. *Microsphaera* spp. [C]
- c. *Oidium* spp. [B]
- d. *Podosphaera* spp. [B]
- e. *Sphaerotheca pannosa* [B]
- f. *Phyllactinia* spp. [B]

5. Rust

- a. *Gymnosporangium juniperi-virginianae* [A]
- b. *Melampsora occidentalis* [A]
- c. *Phragmidium* spp. [B]
- d. *Puccinia* spp. [B]
- e. *Pucciniastrum goeppertianum* [D]
- f. *Uromyces dianthi* [B]

Table 3. Application Regimes

[A] Mix 0.75-1.5 fl. oz. of MPower Propiconazole in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply every 14-21 days during the period of primary infection. If disease is present, tank mix with an EPA-registered contact fungicide. For flower blight, apply MPower Propiconazole when there is 5-10% bloom and again at 70-100% bloom. For dogwoods, apply the 0.75—1.5 fl. oz. rate every 14 days or apply 3 fl. oz. of MPower Propiconazole every 28 days.

[B] Mix 1.8-3.0 fl. oz. of MPower Propiconazole in 100 gallons of water and apply as a full coverage spray to the point of drip. Begin applying when conditions are favorable for disease development. For black spot, apply with a registered contact fungicide labeled for black spot. For Calendula, apply every 30 days. For diplodia tip blight, make 3 applications every 14 days prior to major period of infection. For juniper phomopsis

blight, make the first application as soon as junipers start to grow, and repeat the applications every 14-21 days during periods of active growth.

[C] Mix 3-4.5 fl. oz. of MPower Propiconazole in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply every 30 days, beginning when conditions are favorable for the disease development. For pecans, apply the 4.5 fl. oz. rate beginning at bud break. Apply 3 times at 14-day intervals. For walnuts, apply 3 fl. oz. at 14 to 21 day intervals. For ray blight, apply 4.5 fl. oz. at 7 day intervals or 7.5 fl. oz. at 14 day intervals. For impatiens, bayberry, linden, magnolia, sweet gum and wax myrtle, the maximum use rate is 8 fl. oz.

[D] Mix 6 fl. oz. of MPower Propiconazole in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply every 14-28 days, beginning when conditions are favorable for disease development. For Douglas fir needle rust, apply once in May. For tip blight, start applications in mid-late winter and apply 3 times at 2-month intervals.

[E] Mix 7.5-8.7 fl. oz. of MPower Propiconazole in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply during June to August time period.

Restrictions: Do not apply to apple, Bartlett pear, cherry, citrus, nectarine, peach, pear, pecan, plum or walnut trees that will bear harvestable fruit within 12 months.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in the original container only.

PESTICIDE DISPOSAL: Pesticide wastes may be toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable Container (equal to or less than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling (if available), or dispose of in a sanitary landfill, or by other state and local approved procedures.

Nonrefillable Container (greater than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Return to manufacturer, or offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other state and local approved procedures.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. Use of this product by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

AgraCity Crops & Nutrition, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. AgraCity Crops & Nutrition, Inc. **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of AgraCity Crops & Nutrition, Inc. or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at AgraCity Crops & Nutrition, Inc. election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, AgraCity Crops & Nutrition, Inc. shall not be liable for losses or damages resulting from handling or use of this product. In no case, to the extent consistent with applicable law, shall AgraCity Crops & Nutrition, Inc. be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of AgraCity Crops & Nutrition, Inc. or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

[EPA approval date]